

# Hunting for Quarks: New Physics at Jefferson Lab

*G.P. Gilfoyle*

*University of Richmond, Richmond, VA 23173*

## Outline

- Jefferson Lab's Mission
- What we know.
- What we don't know.
- What we'll learn.
- How we'll do it.
- Concluding Remarks



May 8, 2018

# Hunting for Quarks: New Physics at Jefferson Lab

*G.P. Gilfoyle*

*University of Richmond, Richmond, VA 23173*

## Outline

- Jefferson Lab's Mission
- What we know.
- What we don't know.
- What we'll learn.
- How we'll do it.
- Concluding Remarks



# What is the Mission of Jefferson Lab?

- Basic research into the nature of the nucleus and the nucleon.
- Probe the quark-gluon structure of hadronic matter and how it evolves within nuclei.
- Map the geography of the transition from proton-neutron picture of nuclei to one based on quarks and gluons.
- Test Quantum Chromodynamics (QCD) and quark confinement.
- One of the Millennium Prize Problems (Clay Mathematics Institute).



# What is the Mission of Jefferson Lab?

- Basic research into the nature of the nucleus and the nucleon.
- Probe the quark-gluon structure of hadronic matter and how it evolves within nuclei.
- Map the geography of the transition from proton-neutron picture of nuclei to one based on quarks and gluons.
- Test Quantum Chromodynamics (QCD) and quark confinement.
- One of the Millennium Prize Problems (Clay Mathematics Institute).



Jefferson Lab has completed the 12 GeV Upgrade which doubled the CEBAF accelerator energy.


# What Do We Know?

- The Universe is made of quarks and leptons and the force carriers.

BOSONS			force carriers spin = 0, 1, 2, ...		
Unified Electroweak spin = 1			Strong (color) spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge	Name	Mass GeV/c <sup>2</sup>	Electric charge
$\gamma$ photon	0	0	<b>g</b> gluon	0	0
<b>W<sup>-</sup></b>	80.39	-1	Higgs Boson spin = 0		
<b>W<sup>+</sup></b>	80.39	+1	Name	Mass GeV/c <sup>2</sup>	Electric charge
W bosons			<b>H</b> Higgs	126	0
<b>Z<sup>0</sup></b>	91.188	0			
Z boson					

FERMIONS						matter constituents spin = 1/2, 3/2, 5/2, ...	
Leptons spin = 1/2			Quarks spin = 1/2				
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge		
$\nu_L$ lightest neutrino*	$(0-2) \times 10^{-9}$	0	<b>u</b> up	0.002	2/3		
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3		
$\nu_M$ middle neutrino*	$(0.009-2) \times 10^{-9}$	0	<b>c</b> charm	1.3	2/3		
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3		
$\nu_H$ heaviest neutrino*	$(0.05-2) \times 10^{-9}$	0	<b>t</b> top	173	2/3		
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3		

- The atomic nucleus is made of protons and neutrons bound by the strong force.
- The quarks are confined inside the protons and neutrons.
- Protons and neutrons are NOT confined.




# What Do We Know?

- The Universe is made of quarks and leptons and the force carriers.

BOSONS			force carriers spin = 0, 1, 2, ...		
Unified Electroweak spin = 1			Strong (color) spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge	Name	Mass GeV/c <sup>2</sup>	Electric charge
$\gamma$ photon	0	0	<b>g</b> gluon	0	0
<b>W<sup>-</sup></b>	80.39	-1	Higgs Boson spin = 0		
<b>W<sup>+</sup></b>	80.39	+1	Name	Mass GeV/c <sup>2</sup>	Electric charge
W bosons			<b>H</b> Higgs	126	0
<b>Z<sup>0</sup></b>	91.188	0			
Z boson					

- The atomic nucleus is made of protons and neutrons bound by the strong force.
- The quarks are confined inside the protons and neutrons.
- Protons and neutrons are NOT com...

FERMIONS					matter constituents spin = 1/2, 3/2, 5/2, ...	
Leptons spin = 1/2			Quarks spin = 1/2			
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge	
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3	
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3	
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3	
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3	
$\nu_T$						
$\tau$						



# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.

FERMIONS					
matter constituents spin = 1/2, 3/2, 5/2, ...					
Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.
- A quiz: How much does the proton weigh?

FERMIONS					
Leptons			Quarks		
spin = 1/2			spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.
- A quiz: How much does the proton weigh?

$$m_p = 2m_{up} + m_{down}$$

FERMIONS					
matter constituents spin = 1/2, 3/2, 5/2, ...					
Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.
- A quiz: How much does the proton weigh?

FERMIONS					
matter constituents spin = 1/2, 3/2, 5/2, ...					
Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

$$m_p = 2m_{up} + m_{down} = 2(2 \text{ MeV}/c^2) + 5 \text{ MeV}/c^2$$

# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.
- A quiz: How much does the proton weigh?

FERMIONS					
matter constituents spin = 1/2, 3/2, 5/2, ...					
Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

$$m_p = 2m_{up} + m_{down} = 2(2 \text{ MeV}/c^2) + 5 \text{ MeV}/c^2$$

$$= 939 \text{ MeV}/c^2 \quad \text{OOOPS!!!????}$$

# What Don't We Know?

- Matter comes in pairs of quarks or triplets.
- We are mostly triplets (protons and neutrons).
- More than 99% of our mass is in nucleons.
- Proton  $\rightarrow$  2 ups + 1 down.
- Neutron  $\rightarrow$  1 up + 2 downs.
- A quiz: How much does the proton weigh?

**FERMIONS** matter constituents  
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_L$ lightest neutrino*	$(0-2)\times 10^{-9}$	0	<b>u</b> up	0.002	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.005	-1/3
$\nu_M$ middle neutrino*	$(0.009-2)\times 10^{-9}$	0	<b>c</b> charm	1.3	2/3
$\mu$ muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_H$ heaviest neutrino*	$(0.05-2)\times 10^{-9}$	0	<b>t</b> top	173	2/3
$\tau$ tau	1.777	-1	<b>b</b> bottom	4.2	-1/3

$$m_p = 2m_{up} + m_{down} = 2(2 \text{ MeV}/c^2) + 5 \text{ MeV}/c^2$$

$$= 939 \text{ MeV}/c^2 \quad \text{OOOPS!!!????}$$

- $m_n - m_p = 1.29333205(48) \text{ MeV}/c^2$  (exp) Sz. Borsanyi et al. *Science* 347, 1452 (2015).
- $= 1.51(16)(23) \text{ MeV}/c^2$  (th)

# How Do We Learn What's Inside the Nucleon?

- Nucleon elastic electromagnetic form factors (EEFFs) describe the distribution of charge and magnetization in the nucleon.
- They encode the deviations from point-particle behavior.
- Reveal the internal quark-gluon landscape of the nucleon and nuclei.
- We are in the region where the quarks get dressed.
- Rigorously test QCD in the non-perturbative regime.
- Jargon:  $G_E^p$ ,  $G_M^p$ ,  $G_E^n$ ,  $G_M^n$ .





FIG. 5. Curve (a) shows the theoretical Mott curve for a spinless point proton. Curve (b) shows the theoretical curve for a point proton with the Dirac magnetic moment, curve (c) the theoretical curve for a point proton having the anomalous contribution in addition to the Dirac value of magnetic moment. The theoretical curves (b) and (c) are due to Rosenbluth.<sup>8</sup> The experimental curve falls between curves (b) and (c). This deviation from the theoretical curves represents the effect of a form factor for the proton and indicates structure within the proton, or alternatively, a breakdown of the Coulomb law. The best fit indicates a size of  $0.70 \times 10^{-13}$  cm.

McAllister and Hofstadter, PR 102, 851 (1956)

# How Do We Learn What's Inside the Nucleon?

- Nucleon elastic electromagnetic form factors (EEFFs) describe the distribution of charge and magnetization in the nucleon.
- They encode the deviations from point-particle behavior.
- Reveal the internal quark-gluon landscape of the nucleon and nuclei.
- We are in the region where the quarks get dressed.
- Rigorously test QCD in the non-perturbative regime.
- Jargon:  $G_E^p$ ,  $G_M^p$ ,  $G_E^n$ ,  $G_M^n$ .




Robert Hofstadter, Nobel Prize 1961

the theoretical Mott curve for a spinless particle, curve (b) the theoretical curve for a point particle having the anomalous contribution in the form factor of magnetic moment. The theoretical curve (c) is due to Rosenbluth.<sup>8</sup> The experimental data points are shown in (b) and (c). This deviation from the Mott law is the effect of a form factor for the proton, or alternatively, the effect of a form factor for the neutron. The best fit indicates a size of about 1 fm.

Hofstadter, PR 102, 851 (1956)

# How Do We Learn What's Inside the Nucleon?

- Nucleon elastic electromagnetic form factors (EEFFs) describe the distribution of charge and magnetization in the nucleon.
- They encode the deviations from point-particle behavior.
- Reveal the internal quark-gluon landscape of the nucleon and nuclei.
- We are in the region where the quarks get dressed.
- Rigorously test QCD in the non-perturbative regime.
- Jargon:  $G_E^p$ ,  $G_M^p$ ,  $G_E^n$ ,  $G_M^n$ .



the theoretical Mott curve for a spinless particle, (b) the theoretical curve for a point particle having the anomalous contribution in the form factor of magnetic moment. The theoretical curve due to Rosenbluth.<sup>8</sup> The experimental data points (b) and (c). This deviation from the Mott curve indicates the effect of a form factor for the proton, or alternatively, the finite size of the nucleon.

Hofstadter, PR 102, 851 (1956)




Robert Hofstadter, Nobel Prize 1961

# What is a Form Factor?

- Start with the cross section.

$$\frac{d\sigma}{d\Omega} = \frac{\text{scattered flux/solid angle}}{\text{incident flux/surface area}}$$




For elastic scattering use the Rutherford cross section.

# What is a Form Factor?

- Start with the cross section.

$$\frac{d\sigma}{d\Omega} = \frac{\text{scattered flux/solid angle}}{\text{incident flux/surface area}}$$



For elastic scattering use the Rutherford cross section.


- Get the cross section for elastic scattering by point particles with spin.

$$\frac{d\sigma}{d\Omega} = \frac{Z^2 \alpha^2 (\hbar c)^2}{4E^2 \sin^4(\theta/2)} \left(1 - \beta^2 \sin^2 \frac{\theta}{2}\right) \quad (\text{Mott cross section})$$

# What is a Form Factor?

- Start with the cross section.

$$\frac{d\sigma}{d\Omega} = \frac{\text{scattered flux/solid angle}}{\text{incident flux/surface area}}$$



For elastic scattering use the Rutherford cross section.

- Get the cross section for elastic scattering by point particles with spin.

$$\frac{d\sigma}{d\Omega} = \frac{Z^2\alpha^2(\hbar c)^2}{4E^2\sin^4(\theta/2)} \left(1 - \beta^2\sin^2\frac{\theta}{2}\right) \quad (\text{Mott cross section})$$

- What happens when the beam is electrons and the target is not a point?


$$\frac{d\sigma}{d\Omega} = \frac{Z^2\alpha^2(\hbar c)^2}{4E^2\sin^4(\theta/2)} \left(1 - \beta^2\sin^2\frac{\theta}{2}\right) |F(Q^2)|^2$$

where  $Q^2$  is the 4-momentum transfer.

# What is a Form Factor?

- Start with the cross section.

$$\frac{d\sigma}{d\Omega} = \frac{\text{scattered flux/solid angle}}{\text{incident flux/surface area}}$$



For elastic scattering use the Rutherford cross section.

- Get the cross section for elastic scattering by point particles with spin.

$$\frac{d\sigma}{d\Omega} = \frac{Z^2 \alpha^2 (\hbar c)^2}{4E^2 \sin^4(\theta/2)} \left(1 - \beta^2 \sin^2 \frac{\theta}{2}\right) \quad (\text{Mott cross section})$$

- What happens when the beam is electrons and the target is not a point?

$$\frac{d\sigma}{d\Omega} = \frac{Z^2 \alpha^2 (\hbar c)^2}{4E^2 \sin^4(\theta/2)} \left(1 - \beta^2 \sin^2 \frac{\theta}{2}\right) |F(Q^2)|^2$$

where  $Q^2$  is the 4-momentum transfer.

**THE FORM FACTOR!**

# What Is a Form Factor?

- The chain of reason.

$$\frac{d\sigma}{d\Omega} \rightarrow |F(Q^2)|^2 \Leftrightarrow F(Q^2) \leftarrow \rho(\vec{r}) \leftarrow \psi(\vec{r}) \leftarrow \begin{matrix} \text{QCD,} \\ \text{Constituent quarks} \end{matrix}$$

**Experiment**                      **Comparison**                      **Theory**

The form factors are the meeting ground between theory and experiment.

- The Fourier transform of the form factors are related to the charge and current distributions within the neutron.

# What We'll Learn - The Campaign

## The JLab Lineup

Quantity	Method	Target	$Q^2$ (GeV <sup>2</sup> )	Hall	Beam Days
$G_M^p$ *	Elastic scattering	$LH_2$	7 – 15.5	A	24
$G_E^p/G_M^p$	Polarization transfer	$LH_2$	5 – 12	A	45
$G_M^n$	$E - p/e - n$ ratio	$LD_2, LH_2$	3.5 – 13.0	B	30
$G_M^n$	$E - p/e - n$ ratio	$LD_2, LH_2$	3.5 – 13.5	A	25
$G_E^n/G_M^n$	Double polarization asymmetry	polarized $^3\text{He}$	5 – 8	A	50
$G_E^n/G_M^n$	Polarization transfer	$LD_2$	4 – 7	C	50
$G_E^n/G_M^n$	Polarization transfer	$LD_2$	4.5	A	5

\* Data collection is complete.


PAC approval for 229 days of running in the first five years.

# What We'll Learn - Flavor Decomposition

- With all four EEFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. **194**, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.





# What We'll Learn - Flavor Decomposition

- With all four EEEFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. **194**, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.





- Agreement with Nambu-Jona-Lasinio model encouraging - no parameter fits to the EEEFs.

# What We'll Learn - Flavor Decomposition

- With all four EEEFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. **194**, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.




- Agreement with Nambu-Jona-Lasinio model encouraging - no parameter fits to the EEEFs.

The JLab program will double our reach in  $Q^2$  to  $\approx 8 \text{ GeV}^2$ .

# What We'll Learn - Dyson-Schwinger Eqs


- Equations of motion of quantum field theory.

- Infinite set of coupled integral equations.
- Inherently relativistic, non-perturbative, connected to QCD.
- Deep connection to confinement, dynamical chiral symmetry breaking.
- Infinitely many equations, gauge dependent  $\rightarrow$  Choose well!



- Recent results (Cloët et al).


- Model the nucleon dressed quark propagator as a quark-diquark.
- Damp the shape of the mass function  $M(p)$ .



# What We'll Learn - Dyson-Schwinger Eqs


- Equations of motion of quantum field theory.

- Infinite set of coupled integral equations.
- Inherently relativistic, non-perturbative, connected to QCD.
- Deep connection to confinement, dynamical chiral symmetry breaking.
- Infinitely many equations, gauge dependent  $\rightarrow$  Choose well!



- Recent results (Cloët et al).


- Model the nucleon dressed quark propagator as a quark-diquark.
- Damp the shape of the mass function  $M(p)$ .



# What We'll Learn - Dyson-Schwinger Eqs

- Equations of motion of quantum field theory.


- Infinite set of coupled integral equations.
- Inherently relativistic, non-perturbative, connected to QCD.
- Deep connection to confinement, dynamical chiral symmetry breaking.
- Infinitely many equations, gauge dependent  $\rightarrow$  Choose well!



- Recent results (Cloët et al).


- Model the nucleon dressed quark propagator as a quark-diquark.
- Damp the shape of the mass function  $M(p)$ .

Position of zero in  $\mu_p G_E^p / G_M^p$  and  $\mu_n G_E^n / G_M^n$  sensitive to shape of  $M(p)$ !





# What We'll Learn - Light Front Holographic QCD

- 1 Based on connections between light-front dynamics, it's holographic mapping to anti-de Sitter space, and conformal quantum mechanics.
- 2 Recent paper by Sufian *et al.* (Phys. Rev. D95, 01411 (2017)) included calculations of the electromagnetic form factors that include higher order Fock components  $|qqq\bar{q}\bar{q}\rangle$ .
- 3 Obtain good agreement with all the form factor data with only three parameters, e.g.  $\mu_n G_E^n / G_M^n$ .




# What We'll Learn - Light Front Holographic QCD

- 1 Based on connections between light-front dynamics, it's holographic mapping to anti-de Sitter space, and conformal quantum mechanics.
- 2 Recent paper by Sufian *et al.* (Phys. Rev. D95, 01411 (2017)) included calculations of the electromagnetic form factors that include higher order Fock components  $|qqq\bar{q}\bar{q}\rangle$ .
- 3 Obtain good agreement with all the form factor data with only three parameters, *e.g.*  $\mu_n G_E^n / G_M^n$ .




# How Will We Get There - Jefferson Lab

- Start at your local mile-long, high-precision, 12-GeV electron accelerator.
- The Continuous Electron Beam Accelerator Facility (CEBAF) produces beams of unrivaled quality.
- Electrons do up to five laps, are extracted, and sent to one of three experimental halls.
- All four halls can run simultaneously.




# How Will We Get There - Jefferson Lab

- Start at your local mile-long, high-precision, 12-GeV electron accelerator.
- The Continuous Electron Beam Accelerator Facility (CEBAF) produces beams of unrivaled quality.
- Electrons do up to five laps, are extracted, and sent to one of three experimental halls.
- All four halls can run simultaneously.





# How Will We Get There - Jefferson Lab

- Start at your local mile-long, high-precision, 12-GeV electron accelerator.
- The Continuous Electron Beam Accelerator Facility (CEBAF) produces beams of unrivaled quality.
- Electrons do up to five laps, are extracted, and sent to one of three experimental halls.
- All four halls can run simultaneously.




# How Will We Get There - Jefferson Lab

- Start at your local mile-long, high-precision, 12-GeV electron accelerator.
- The Continuous Electron Beam Accelerator Facility (CEBAF) produces beams of unrivaled quality.
- Electrons do up to five laps, are extracted, and sent to one of three experimental halls.
- All four halls can run simultaneously.





# How Do We Measure The Form Factors

- Add one 45-ton, \$80-million radiation detector: the CEBAF Large Acceptance Spectrometer (CLAS12).
- CLAS covers a large fraction of the total solid angle at forward angles.
- Has about 62,000 detecting elements in about 40 layers.




# How Do We Measure The Form Factors

- Add one 45-ton, \$80-million radiation detector: the CEBAF Large Acceptance Spectrometer (CLAS12).




# How Do We Measure The Form Factors


- Add one 45-ton, \$80-million radiation detector: the CEBAF Large Acceptance Spectrometer (CLAS12).



# A CLAS12 Event - Summary



# A CLAS12 Event - Summary



# How Do We Extract the Form Factors? - $G_M^n$



- E12-07-104 in Hall B (Gilfoyle, Hafidi, Brooks).
- Ratio Method on Deuterium:

$$R = \frac{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' n)_{QE}]}{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' p)_{QE}]}$$

$$= a \times \frac{\sigma_{\text{Mott}} \left( \frac{(G_E^n)^2 + \tau (G_M^n)^2}{1 + \tau} + 2\tau \tan^2 \frac{\theta_e}{2} (G_M^n)^2 \right)}{\frac{d\sigma}{d\Omega} [{}^1\text{H}(e, e' p)]}$$

where  $a$  is nuclear correction.

- Precise neutron detection efficiency needed to keep systematics low.
  - tagged neutrons from  ${}^2\text{H}(e, e' pn)$ .
  - $\text{LH}_2$  target.
- Kinematics:  $Q^2 = 3.5 - 13.0 \text{ (GeV}/c)^2$ .
- Beamtime: 40 days.
- Systematic uncertainties  $< 2.5\%$  across full  $Q^2$  range.
- Run Group B starts February, 2019(?).



# How Do We Extract the Form Factors? - $G_M^n$



- E12-07-104 in Hall B (Gilfoyle, Hafidi, Brooks).
- Ratio Method on Deuterium:

$$R = \frac{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' n)_{QE}]}{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' p)_{QE}]}$$

$$= a \times \frac{\sigma_{\text{Mott}} \left( \frac{(G_E^n)^2 + \tau (G_M^n)^2}{1 + \tau} + 2\tau \tan^2 \frac{\theta_e}{2} (G_M^n)^2 \right)}{\frac{d\sigma}{d\Omega} [{}^1\text{H}(e, e' p)]}$$


where  $a$  is nuclear correction.

- Precise neutron detection efficiency needed to keep systematics low.
  - tagged neutrons from  ${}^2\text{H}(e, e' pn)$ .
  - $LH_2$  target.
- Kinematics:  $Q^2 = 3.5 - 13.0 \text{ (GeV}/c)^2$ .
- Beamtime: 40 days.
- Systematic uncertainties  $< 2.5\%$  across full  $Q^2$  range.
- Run Group B starts February, 2019(?).



# Concluding Remarks

- JLab is a laboratory to test and expand our understanding of quarks, gluons, nuclear matter and QCD.
- We continue to unravel the nature of matter at greater and greater depths.
- Lots of new and exciting results are coming out.
- A bright future lies ahead in the 12 GeV Era.



# Some Facts of Life On The Frontier

- Work at Jefferson Lab in Newport News.
  - 700 physicists, engineers, technicians, and staff.
  - Vibrant intellectual environment - talks, visitors, educational programs...
  - Lots going on.
- Richmond group part of CLAS Collaboration.
  - operates CLAS12.
  - ~190 physicists, 40 institutions, 13 countries.
  - Part of Software Group - emphasis on software development.
  - Past Surrey masters students (and Richmond undergrads) have presented posters at meetings, appeared on JLab publications,....
- Run-Group B consists of seven experiments (including  $G_M^n$ ) and is expected to run in spring 2019.



# Additional Slides

# Some Necessary Background

- EEFs cross section described with Dirac ( $F_1$ ) and Pauli ( $F_2$ ) form factors

$$\frac{d\sigma}{d\Omega} = \sigma_{Mott} \left[ (F_1^2 + \kappa^2 \tau F_2^2) + 2\tau (F_1 + \kappa F_2)^2 \tan^2 \left( \frac{\theta_e}{2} \right) \right]$$

where

$$\sigma_{Mott} = \frac{\alpha^2 E' \cos^2(\frac{\theta_e}{2})}{4E^3 \sin^4(\frac{\theta_e}{2})}$$

and  $\kappa$  is the anomalous magnetic moment,  $E$  ( $E'$ ) is the incoming (outgoing) electron energy,  $\theta$  is the scattered electron angle and  $\tau = Q^2/4M^2$ .

- For convenience use the Sachs form factors.





$$\frac{d\sigma}{d\Omega} = \frac{\sigma_{Mott}}{\epsilon(1+\tau)} (\epsilon G_E^2 + \tau G_M^2)$$

where


$$G_E = F_1 - \tau F_2 \quad \text{and} \quad G_M = F_1 + F_2 \quad \text{and} \quad \epsilon = \left[ 1 + 2(1 + \tau) \tan^2 \frac{\theta_e}{2} \right]^{-1}$$

# Where We Are Now.


- $G_M^p$  well known over large  $Q^2$  range.
- The ratio  $G_E^p/G_M^p$  from polarization transfer measurements diverged from previous Rosenbluth separations.
  - Two-photon exchange (TPE).
  - Effect of radiative corrections.
- Neutron magnetic FF  $G_M^n$  still follows dipole.
- High- $Q^2$   $G_E^n$  opens up flavor decomposition.




# The Experiments - New Detectors



Hall A - High Resolution Spectrometer (HRS) pair, SuperBigBite (SBS), neutron detector, and specialized installation experiments.




Hall B - CLAS12 large acceptance spectrometer operating at high luminosity with toroid (forward detector) and solenoid (central detector).



Hall C - New Super High Momentum Spectrometer to paired with the existing High Momentum Spectrometer.

Hall D - A new large acceptance detector based on a solenoid magnet for photon beams is under construction.

## GlueX Detector



# Extracting $G_M^n$

- Use ratio method on deuterium:

$$R = \frac{\frac{d\sigma}{d\Omega}[{}^2\text{H}(e,e'n)_{QE}]}{\frac{d\sigma}{d\Omega}[{}^2\text{H}(e,e'p)_{QE}]} = a \times \frac{\sigma_{\text{Mott}} \left( \frac{(G_E^n)^2 + \tau(G_M^n)^2}{1+\tau} + 2\tau \tan^2 \frac{\theta_e}{2} (G_M^n)^2 \right)}{\frac{d\sigma}{d\Omega}[{}^1\text{H}(e,e')p]}$$



where  $a$  is a nuclear correction.

- Acceptance matching on  $e - p$  and  $e - n$  measurements. For each event swim both nucleons through CLAS12 and require both to strike the CLAS12 fiducial volume it to be accepted.
- Select quasi-elastic events by requiring the nucleon scattering angle to be within a narrow angular cone around the direction predicted by elastic scattering (no Fermi motion).
- Require no other particles in the final state to reduce inelastic contributions.
- Apply neutron/proton detection efficiency, Fermi motion, nuclear corrections and others to  $R$ .

# Neutron Magnetic Form Factor $G_M^n$ - 2

- E12-09-019 in Hall A (Quinn, Wojtsekhowski, Gilman).
- Ratio Method on Deuterium as in Hall B:  

$$R = \frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' n)_{QE}] / \frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' p)_{QE}]$$
- Electron arm: SuperBigBite spectrometer.
- Hadron arm: hadron calorimeter (HCal).
- Neutron detection efficiency:
  - Use  $p(\gamma, \pi^+)n$  for tagged neutrons.
  - End-point method.
- Kinematics:  $Q^2 = 3.5 - 13.5 \text{ (GeV}/c)^2$ .
- Beamtime: 25 days.
- Systematic uncertainties  $< 2.1\%$ .
- Two  $G_M^n$  measurements 'allow a better control for the systematic error' (PAC34).
- Expected in next 2-3 years.





# Proton Magnetic Form Factor - $G_M^p$

- E12-07-108 in Hall A (Gilad, Moffitt, Wojtsekhowski, Arrington).
- Precise measurement of  $ep$  elastic cross section and extract  $G_M^p$ .
- Both HRSs in electron mode.
- Beamtime: 24 days.
- $Q^2 = 7.0 - 15.5 \text{ GeV}^2$  (1.0, 1.5  $\text{GeV}^2$  steps).
- Significant reduction in uncertainties:

	$d\sigma/d\Omega$	$G_M^p$
Point-to-Point	1.0-1.3	0.5-0.6
Normalization	1.0-1.3	0.5-0.6
Theory	1.0-2.0	0.5-1.0

- Two-Photon Exchange is a major source of uncertainty  $\rightarrow$  vary  $\epsilon$  to constrain.
- Sets the scale of other EEEFs.
- Completed data collection in 2017.



E. Christy, Hall A Summer Meeting 2017