Luminosity dependence of $N_{eh\pm}/N_e$, RG-A Data 1



CLAS12-NOTE 2020-005



S.Stepanyan, RG-B, 9/7/24

Luminosity dependence of RGB data 2

RGB Pass 2 Review



Normalized yields of eh+/-

Conventional
AI-assisted
Ratio

RGB N_{ep}/N_e luminosity dependence with QE cuts on - beam energy calculated from particle angles, θ_{pq}, and $\Delta\phi$.





RGB N_{ep}/N_e luminosity dependence with QE cuts off - dropped cuts on beam energy calculated from particle angles, θ_{pq}, and $\Delta\phi$.





RGB N_{ep}/N_e luminosity dependence with QE cuts on/off.



Results from four data sets 6









Comparison of Tracking Efficiency Slopes 7

	QE cuts on	QE cuts off	$\mathrm{E_{beam}[GeV]}$
Spring, 2020	$-0.0076 \pm 0.0023\ nA^{-1}$	$-0.0082 \pm 0.0001\ nA^{-1}$	10.4
Fall, 2019	$-0.0104 \pm 0.0021\ nA^{-1}$	$-0.0079 \pm 0.0001\ nA^{-1}$	10.4
Spring, 2019	$-0.0075 \pm 0.0018\ nA^{-1}$	$-0.0087 \pm 0.0001\ nA^{-1}$	10.2
Spring, 2019	$-0.0093 \pm 0.0013\ nA^{-1}$	$-0.0082 \pm 0.0001\ nA^{-1}$	10.6

Weighted averages:

For QE cuts on: $\langle b \rangle = 0.0087 \pm 0.0014\ nA^{-1}$

For QE cuts off: $\langle b \rangle = 0.00883 \pm 0.00005\ nA^{-1}$

Extracting average beam currents 8

1. Use NB method to get the current at frequent intervals - Get ungated Faraday cut reading in RUN::config bank which records integrated current since last time the bank was written out.
2. Get the timestamp from Trigger Interface board in RUN::config.
3. Divide by the time since the last recording of RUN::config bank.
4. Require the current exceed a threshold of 1 nA to remove beam trips.

From CLAS12-NOTE 2020-005

1. $p_{\pm} > 0.4\ \mathrm{GeV/c}$
2. $|\chi 2PID| < 5$, $|\chi 2| < 10|$
3. Reject tracks with FTOF Panel 2 - don't see any.
4. Vertex $-15 < v_z < 5\ cm$ in RGA. RGB had $-13 < v_z < 12\ cm$ for inbending.
5. fiducial cuts