

## Magnetic Structure of the Ground State Neutron

The electromagnetic form factors of the proton and neutron are fundamental observables that describe the internal distribution of charge and electric current. The electromagnetic structure of the proton ground state has been mapped out for over 50 years and with high precision and in a large range of  $Q^2$ . However, only recently has it become possible to access the magnetic structure of the neutron with similarly high precision. A result [1] from the CLAS collaboration measured the magnetic form factor of the neutron using the ratio method on deuterium (right-hand panel, red points in figure). The precision and coverage of these results eclipse the world's data on this elastic form factor in the  $Q^2$  range of 1.0 – 4.8  $\text{GeV}^2$  (green points). The experiment used a double-cell target made from liquid hydrogen and liquid deuterium and detected the quasi-elastic reactions  $en(p_s) \rightarrow en$  and  $ep(n_s) \rightarrow ep$  from the deuterium cell and  $ep \rightarrow e\pi^+n$  from the hydrogen target. The first two quasi-elastic reactions measure the cross section ratio of the neutron and proton simultaneously, which allows the control of systematic uncertainties at the 2-3% level. The  $\pi^+n$  reaction on hydrogen allows a precise calibration of the neutron detection efficiency *in situ* and simultaneous with the physics reaction. The left panel in the figure shows the neutron detection efficiencies for the electromagnetic calorimeters and the time-of-flight plastic scintillators. Surprisingly, the CLAS6 result deviates from the dipole form by not more than 10% in the entire  $Q^2$  range covered (see theory curves in the figure). An experiment to push this measurement to nearly three times the upper  $Q^2$  limit of the current CLAS6 data has been approved for running in Hall B with the new CLAS12 detector after the 12-GeV Upgrade is completed. The anticipated data range and precision is shown in the right panel (black points).




Figure: The left panel shows the results of the neutron detection efficiency for the CLAS6 electromagnetic calorimeters and the TOF system (inset). The right panel shows the magnetic form factor of the neutron normalized to the dipole form factor with results from other measurements and theoretical curves.

[1] J. Lachniet *et al.* (CLAS collaboration), Phys. Rev. Lett. **102**, 192001 (2009).