

---

# Future Measurements of the Nucleon Elastic Electromagnetic Form Factors at Jefferson Lab

*G.P. Gilfoyle*

*University of Richmond, Richmond, VA 23173*

## Outline

1. Scientific Motivation
2. Necessary Background
3. What we hope to learn.
4. The Measurements
5. Summary and Conclusions



Iglesia de la Matriz

# Scientific Motivation - What We Hope to Learn.

---

- Nucleon elastic electromagnetic form factors (EEFFs) describe the distribution of charge and magnetization in the nucleon.
- Reveal the internal landscape of the nucleon and nuclei.
- Rigorously test QCD in the non-perturbative regime.
  - Nuclear models, constituent quarks,...
  - lattice QCD.
- Map the transition from the hadronic picture to QCD.

# Scientific Motivation - What We Hope to Learn.

---

- Nucleon elastic electromagnetic form factors (EEFFs) describe the distribution of charge and magnetization in the nucleon.
- Reveal the internal landscape of the nucleon and nuclei.
- Rigorously test QCD in the non-perturbative regime.
  - Nuclear models, constituent quarks,...
  - lattice QCD.
- Map the transition from the hadronic picture to QCD.

EEFFs have played an essential role in nuclear and nucleon structure for more than a half century.

# Some Necessary Background

- EEFFs cross section described with Dirac ( $F_1$ ) and Pauli ( $F_2$ ) form factors

$$\frac{d\sigma}{d\Omega} = \sigma_{Mott} \left[ (F_1^2 + \kappa^2 \tau F_2^2) + 2\tau (F_1 + \kappa F_2)^2 \tan^2 \left( \frac{\theta_e}{2} \right) \right]$$

where

$$\sigma_{Mott} = \frac{\alpha^2 E' \cos^2 \left( \frac{\theta_e}{2} \right)}{4E^3 \sin^4 \left( \frac{\theta_e}{2} \right)}$$

and  $\kappa$  is the anomalous magnetic moment,  $E$  ( $E'$ ) is the incoming (outgoing) electron energy,  $\theta$  is the scattered electron angle and  $\tau = Q^2/4M^2$ .

- For convenience use the Sachs form factors.

$$\frac{d\sigma}{d\Omega} = \sigma_{Mott} \left( \frac{(G_E^n)^2 + \tau(G_M^n)^2}{1 + \tau} + 2\tau \tan^2 \frac{\theta_e}{2} (G_M^n)^2 \right)$$


where

$$G_E = F_1 - \tau F_2$$


$$G_M = F_1 + F_2$$

# Where We Are Now.


- The ratio  $G_E^p/G_M^p$  from recoil polarization measurements diverged from previous Rosenbluth separations.
  - Two-photon exchange (TPE).
  - Effect of radiative corrections.
- Neutron magnetic FF  $G_M^n$  still follows dipole.
- High- $Q^2$   $G_E^n$  opens the door to flavor decomposition.



PRL 104, 242301 (2010)




PRL 105, 262302 (2010)




Scholarpedia, 5(8):10204

# Where We Are Now.

- The ratio  $G_E^p/G_M^p$  from recoil polarization measurements diverged from previous Rosenbluth separations.
  - Two-photon exchange (TPE).
  - Effect of radiative corrections.
- Neutron magnetic FF  $G_M^n$  still follows dipole
- High- $Q^2$   $G_E^n$  opens the door




PRL 104, 242301 (2010)



PRL 105, 262302 (2010)

Advances driven by:

- high luminosity beams
- large acceptance detectors
- polarized beams, targets, detectors




Scholarpedia, 5(8):10204

# Where We Are Now.

The EEFFs emerge from Quantum Chromodynamics (QCD), but calculations here require non-perturbative methods.



- Vector Meson Dominance and dispersion analyses fit all four EEFFs, but use many parameters.
- Constituent Quark Models highlight relativity, but don't capture all of QCD.
- Generalized Parton Distributions (GPDs) connect valence quarks in transverse space and longitudinal momentum.
- EEFFs are the first moments of the GPDs.



# Where We Are Now.

The EEFFs emerge from Quantum Chromodynamics (QCD), but calculations here require non-perturbative methods.

- Vector Meson Dominance and dispersion analyses fit all four EEFFs, but use many parameters.
- Constituent Quark Models highlight relativity, but don't capture all of QCD.
- Generalized Parton Distributions (GPDs) connect valence quarks in transverse space and longitudinal momentum.
- EEFFs are the first moments of the GPDs.





# Where We Are Now - Lattice QCD

- Lattice gauge theory is the only means of *ab initio* QCD calculations in the non-perturbative regime.
- Computationally challenging.
- EEFFs are an early test of IQCD.
  - The isovector form of the EEFFs is


$$F_{1,2}^V = \frac{F_{1,2}^p - F_{1,2}^n}{2}$$

where

$$F_1 = \frac{\tau G_M + G_E}{1 + \tau} \quad F_2 = \frac{G_M - G_E}{1 + \tau}$$


where  $\tau = Q^2/4M^2$ .

- This form of the EEFFs does not have disconnected diagrams which are computationally intensive.
- Expect EEFF calculation in the next decade.




# Where We Are Going - Dyson-Schwinger Eqs

- Equations of motion of quantum field theory.
  - Infinite set of coupled integral equations.
  - Inherently relativistic, non-perturbative, connected to QCD.
  - Deep connection to confinement, dynamical chiral symmetry breaking.
  - Infinitely many equations, gauge dependent → Choose well!
- Recent results (Cloët et al).
  - Model the nucleon dressed quark propagator as a quark-diquark.
  - Damp the shape of the mass function  $M(p)$ .




# Where We Are Going - Dyson-Schwinger Eqs

- Equations of motion of quantum field theory.
  - Infinite set of coupled integral equations.
  - Inherently relativistic, non-perturbative, connected to QCD.
  - Deep connection to confinement, dynamical chiral symmetry breaking.
  - Infinitely many equations, gauge dependent → Choose well!



- Recent results (Cloët et al).
  - Model the nucleon dressed quark propagator as a quark-diquark.
  - Damp the shape of the mass function  $M(p)$ .


Position of zero in  $\mu_p G_{Ep}^p / G_M^p$   
sensitive to shape of  $M(p)$ !



# Where We Are Going - Flavor Decomposition

- With all four EEFFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. **194**, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$




PRL **106**, 252003 (2011).

# Where We Are Going - Flavor Decomposition

- With all four EEFFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. **194**, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.




PRL **106**, 252003 (2011).

# Where We Are Going - Flavor Decomposition


- With all four EEFFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. 194, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.



PRC, 86 065210



PRL 106, 252003 (2011).



- $F_2^f / F_2^u$  ratio not well reproduced by any models  $\rightarrow$  good test bench.

# Where We Are Going - Flavor Decomposition

- With all four EEFFs we can unravel the contributions of the  $u$  and  $d$  quarks.
- Assume charge symmetry, no  $s$  quarks and use (Miller *et al.* Phys. Rep. 194, 1 (1990))

$$F_{1(2)}^u = 2F_{1(2)}^p + F_{1(2)}^n \quad F_{1(2)}^d = 2F_{1(2)}^n + F_{1(2)}^p$$

- Evidence of di-quarks?  $d$ -quark scattering probes the diquark.



PRL 106, 252003 (2011).

- $F_2^f / F_2^u$  ratio not well reproduced by any models  $\rightarrow$  good test bench.

The JLab program will double our reach in  $Q^2$  here.

# Where We Are Going - New Experiments.

## The JLab Lineup



Quantity	Method	Target	$Q^2$ (GeV <sup>2</sup> )	Hall	Beam Days
$G_M^p$	Elastic scattering	$LH_2$	7 – 15.5	A	24
$G_E^p / G_M^p$	Polarization transfer	$LH_2$	5 – 12	A	45
$G_M^n$	$E - p/e - n$ ratio	$LD_2 - LH_2$	3.5 – 13.0	B	30
$G_M^n$	$E - p/e - n$ ratio	$LD_2, LH_2$	3.5 – 13.5	A	25
$G_E^n / G_M^n$	Double polarization asymmetry	polarized $^3\text{He}$	5 – 8	A	50
$G_E^n / G_M^n$	Polarization transfer	$LD_2$	4 – 7	C	50

All are extensions of 6 GeV era experiments.

PAC approval for 224 days of running in the first five years.



# How We Will Get There: Jefferson Lab.



Continuous Electron Beam Accelerator Facility (CEBAF)



- Superconducting Electron Accelerator, 100% duty cycle.
- $E_{max} = 11 \text{ GeV}$  (Halls A, B, and C) and 12 GeV (Hall D),  $\Delta E/E \approx 2 \times 10^{-4}$ ,  $I_{summed} \approx 90 \mu\text{A}$ ,  $P_e \geq 80\%$ .

# Proton Magnetic Form Factor - $G_M^p$

- E12-07-108 in Hall A (Gilad, Moffitt, Wojtsekhowski, Arrington).
- Precise measurement of  $ep$  elastic cross section and extract  $G_M^p$ .
- Both HRSs in electron mode.
- Beamtime: 24 days.
- $Q^2 = 7.0 - 15.5 \text{ GeV}^2$  (1.0, 1.5  $\text{GeV}^2$  steps).
- Significant reduction in uncertainties:

	$d\sigma/d\Omega$	$G_M^p$
Point-to-Point	1.0-1.3	0.5-0.6
Normalization	1.0-1.3	0.5-0.6
Theory	1.0-2.0	0.5-1.0

- Two-Photon Exchange is a major source of uncertainty  $\rightarrow$  vary  $\epsilon$  to constrain.
- Sets the scale of other EEFFs.



# Proton Form Factor Ratio $G_E^p/G_M^p$

- E12-07-109 (GEp(5)) in Hall A (Brash, Jones, Perdrisat, Pentchev, Cisbani, Punjabi, Khandaker, Wojtsekhowski).


- Polarization transfer using  $H(\vec{e}, e' \vec{p})$ :

$$\frac{G_E^p}{G_M^p} = -\frac{P_t}{P_l} \frac{E + E'}{2M} \tan\left(\frac{\theta_e}{2}\right)$$

- Electron arm: EM calorimeter (BigCal).
- Proton arm: new, large-acceptance magnetic spectrometer (SBS) with double polarimeter, and hadron calorimeter.
- Beamtime: 45 days.
- Kinematics and Uncertainties:

$Q^2$ (GeV <sup>2</sup> )	5.0	8.0	12.0
$\Delta[\mu G_E/G_m]$	0.025	0.031	0.069

- Combined with GEp(4).



# Proton Form Factor Ratio $G_E^p/G_M^p$

- E12-07-109 (GEp(5)) in Hall A (Brash, Jones, Perdrisat, Pentchev, Cisbani, Punjabi, Khandaker, Wojtsekhowski).


- Polarization transfer using  $H(\vec{e}, e' \vec{p})$ :

$$\frac{G_E^p}{G_M^p} = -\frac{P_t}{P_l} \frac{E + E'}{2M} \tan\left(\frac{\theta_e}{2}\right)$$

- Electron arm: EM calorimeter (BigCal).
- Proton arm: new, large-acceptance magnetic spectrometer (SBS) with double polarimeter, and hadron calorimeter.
- Beamtime: 45 days.
- Kinematics and Uncertainties:

$Q^2$ (GeV <sup>2</sup> )	5.0	8.0	12.0
$\Delta[\mu G_E/G_m]$	0.025	0.031	0.069

- Combined with GEp(4).





# Proton Form Factor Ratio $G_E^p/G_M^p$

- E12-07-109 (GEp(5)) in Hall A (Brash, Jones, Perdrisat, Pentchev, Cisbani, Punjabi, Khandaker, Wojtsekhowski).

- Polarization transfer using  $H(\vec{e}, e' \vec{p})$ :



$$\frac{G_E^p}{G_M^p} = -\frac{P_t}{P_l} \frac{E + E'}{2M} \tan\left(\frac{\theta_e}{2}\right)$$

- Electron arm: EM calorimeter (BigCal).
- Proton arm: new, large-acceptance magnetic spectrometer (SBS) with double polarimeter, and hadron calorimeter.
- Beamtime: 45 days.
- Kinematics and Uncertainties:

$Q^2$ (GeV <sup>2</sup> )	5.0	8.0	12.0
$\Delta[\mu G_E/G_m]$	0.025	0.031	0.069

- Combined with GEp(4).

Proton form factors ratio, GEp(5) (E12-07-109)



# Neutron Magnetic Form Factor $G_M^n$ - 1



- E12-07-104 in Hall B (Gilfoyle, Hafidi, Brooks).
- Ratio Method on Deuterium ( ${}^2\text{H}(e, e'p)n$  and  ${}^2\text{H}(e, e'n)p$ ):

$$R = \frac{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e'n)_{QE}]}{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e'p)_{QE}]}$$

$$= a \times \frac{\sigma_{Mott} \left( \frac{(G_E^n)^2 + \tau(G_M^n)^2}{1 + \tau} + 2\tau \tan^2 \frac{\theta_e}{2} (G_M^n)^2 \right)}{\frac{d\sigma}{d\Omega} [{}^1\text{H}(e, e')p]}$$

where  $a$  is nuclear correction.

- Precise neutron detection efficiency needed to keep systematics low.
  - tagged neutrons from  $p(e, e'\pi^+n)$ .
  - Dual  $LD_2 - LH_2$  target for *in situ* calibrations.
- Kinematics:  $Q^2 = 3.5 - 13.0 \text{ (GeV}/c)^2$ .
- Beamtime: 30 days.
- Systematic uncertainties less than 2.5% across full  $Q^2$  range.



2011-09-19 11:03:40

# Neutron Magnetic Form Factor $G_M^n$ - 2

- E12-09-019 in Hall A (Quinn, Wojtsekhowski, Gilman).

- Ratio Method on Deuterium as in Hall B:

$$R = \frac{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' n)_{QE}]}{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' p)_{QE}]}$$

- Electron arm: SuperBigBite spectrometer.
- Hadron arm: hadron calorimeter (HCal).
- Neutron detection efficiency:



- Use  $p(\gamma, \pi^+)n$  for tagged neutrons.
- End-point method.

- Kinematics:  $Q^2 = 3.5 - 13.5 \text{ (GeV/c)}^2$ .

- Beamtime: 25 days.

- Systematic uncertainties  $< 2.1\%$ .

- Two  $G_M^n$  measurements 'allow a better control for the systematic error' (PAC34).



# Neutron Magnetic Form Factor $G_M^n$ - 2

- E12-09-019 in Hall A (Quinn, Wojtsekhowski, Gilman).

- Ratio Method on Deuterium as in Hall B:

$$R = \frac{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' n)_{QE}]}{\frac{d\sigma}{d\Omega} [{}^2\text{H}(e, e' p)_{QE}]}$$

- Electron arm: SuperBigBite spectrometer.

- Hadron arm: hadron calorimeter (HCal).

- Neutron detection efficiency:



- Use  $p(\gamma, \pi^+)n$  for tagged neutrons.
- End-point method.

- Kinematics:  $Q^2 = 3.5 - 13.5 \text{ (GeV/c)}^2$ .

- Beamtime: 25 days.

- Systematic uncertainties  $< 2.1\%$ .

- Two  $G_M^n$  measurements 'allow a better control for the systematic error' (PAC34).





# Neutron Form Factor Ratio $G_E^n/G_M^n - 1$

- E12-09-016 in Hall A (Cates, Wojtsekhowski, Riordan).

- Double Polarization Asymmetry:  
Get  $A_{en}^V$  from  ${}^3\text{He}(\vec{e}, e'\vec{n})pp$ .

- Longitudinally polarized electron beam.

- ${}^3\text{He}$  target polarized perpendicular to the momentum transfer.

- Electron arm: SuperBigBite.

- Neutron arm: hadron calorimeter HCal (overlap with GEp(5) and Hall A  $G_M^n$ ).


- Beamtime: 50 days.

- Kinematics and Uncertainties:

$Q^2$ (GeV <sup>2</sup> )	5.0	6.8	8.0
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{stat}$	0.027	0.022	0.032
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{syst}$	0.018	0.021	0.013

$$A_{en}^V = \frac{-2\sqrt{\tau(\tau+1)} \tan(\theta_e/2) \cos \phi^* \sin \theta^* G_E^n/G_M^n}{(G_E^n/G_M^n)^2 + \tau/\epsilon} + \frac{-2\tau\sqrt{1+\tau+(\tau+1)^2 \tan^2(\theta_e/2)} \tan(\theta_e/2) \cos \theta^*}{(G_E^n/G_M^n)^2 + \tau/\epsilon}$$


$$\text{where } \epsilon = 1 / \left( 1 + 2(1 + \tau) \tan^2\left(\frac{\theta_e}{2}\right) \right)$$



# Neutron Form Factor Ratio $G_E^n / G_M^n - 1$

- E12-09-016 in Hall A (Cates, Wojtsekhowski, Riordan).
- Double Polarization Asymmetry: Get  $A_{en}^V$  from  ${}^3\text{He}(\vec{e}, e'n)pp$ .
- Longitudinally polarized electron beam.
- ${}^3\text{He}$  target polarized perpendicular to the momentum transfer.
- Electron arm: SuperBigBite.
- Neutron arm: hadron calorimeter HCal (overlap with GEp(5) and Hall A  $G_M^n$ ).
- Beamtime: 50 days.
- Kinematics and Uncertainties:



$Q^2$ (GeV <sup>2</sup> )	5.0	6.8	8.0
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{stat}$	0.027	0.022	0.032
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{syst}$	0.018	0.021	0.013



# Neutron Form Factor Ratio $G_E^n/G_M^n - 1$

- E12-09-016 in Hall A (Cates, Wojtsekhowski, Riordan).
- Double Polarization Asymmetry: Get  $A_{en}^V$  from  ${}^3\text{He}(\vec{e}, e'n)pp$ .
- Longitudinally polarized electron beam.
- ${}^3\text{He}$  target polarized perpendicular to the momentum transfer.
- Electron arm: SuperBigBite.
- Neutron arm: hadron calorimeter HCal (overlap with GEp(5) and Hall A  $G_M^n$ ).
- Beamtime: 50 days.
- Kinematics and Uncertainties:

$Q^2$ ( $\text{GeV}^2$ )	5.0	6.8	8.0
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{stat}$	0.027	0.022	0.032
$\Delta \left[ \frac{\mu G_E}{G_M} \right]_{syst}$	0.018	0.021	0.013



# Neutron Form Factor Ratio $G_E^n/G_M^n$ - 2

- E12-11-009 in Hall C (Anderson, Arrington, Kowalski, Madey, Plaster, Semenov).

- Polarization transfer using  ${}^2\text{H}(\vec{e}, e'\vec{n})p$ :

$$\frac{G_E^n}{G_M^n} = -\frac{P_t}{P_l} \frac{E + E'}{2M} \tan\left(\frac{\theta_e}{2}\right)$$

- Electron arm: Super High Momentum Spectrometer (SHMS).

- Neutron arm: neutron polarimeter with tapered-gap neutron-spin-precession magnet and proton recoil detection.



- Kinematics:  $Q^2 = 3.95, 6.88$  ( $\text{GeV}/c$ )<sup>2</sup>.

- Beamtime: 50 days.

- Systematic uncertainties about 2-3%.

- Statistical uncertainties about 10-16%.

- Complementary to the  ${}^3\text{He}$  experiment.



# Summary and Conclusions

---

- Large gains over the last decade or so in physics understanding of the EEFFs built on new technologies and capabilities.
- Major changes in our understanding of nucleon structure.
- Jefferson Lab will mount a broad assault on the EEFFs and will significantly expand the physics reach of our understanding.
- Discovery potential in mapping out nucleon structure and understanding QCD.


---

# Additional Slides







# The Experiments - New Detectors



Hall A - High Resolution Spectrometer (HRS) pair, SuperBigBite, neutron detector, and specialized installation experiments.




Hall C - New Super High Momentum Spectrometer to be used with the existing High Momentum Spectrometer.




Hall B - CLAS12 large acceptance spectrometer operating at high luminosity with toroid (forward detector) and solenoid (central detector).

Hall D - A new large acceptance detector based on a solenoid magnet for photon beam experiments is under construction.





# Current World Data on EEFs



J.J.Kelly, Phys. Rev.C, 068202, 2004.

- Proton form factors have small uncertainties and reach higher  $Q^2$ .
- Neutron form factors are sparse and have large uncertainties.
- Significant deviations from the dipole form factor.


# Where We Are Now - Nucleon Structure and GPDs

- Generalized Parton Distributions (GPDs) connect the valence quark distributions in transverse space and longitudinal momentum.
- EEFFs are the first moments of the GPDs and provide an important constraint.


$$\int dx \sum H^q(x, \zeta, t) = F_1(t) \quad \text{Dirac FF}$$

$$\int dx \sum E^q(x, \zeta, t) = F_2(t) \quad \text{Pauli FF}$$


- Unravel the mass  $M(t)$ , angular momentum  $J(t)$ , and force and pressure  $d_1(t)$ .
- Nucleon form factor measurements complement the Semi-Inclusive Deep Inelastic Scattering program.



Transverse spatial distributions.




Longitudinal momentum distributions.




Correlated spatial and momentum distributions.

# Low- $Q^2$ Comparison of $G_E^p/G_M^p$ Data



# CLAS12 Detector and $G_M^n$ Target



luminosity  $\approx 10^{35} \text{ s}^{-1} \text{ cm}^{-2}$ .



Forward Detector	
Angular Range	$5^\circ - 40^\circ$ (outbenders)
$\Delta p/p$	$< 0.01$ @ 5 GeV/c
$\Delta\theta$ (mr)	$< 1$ $p > 2.5$ GeV/c
$\Delta\phi$ (mr)	$< 3$ $p > 2.5$ GeV/c
Neutron Detection	$N_{eff} = 0.1 - 0.6$

# CLAS12 Detector and $G_M^n$ Target

luminosity  $\approx 10^{35} \text{ s}^{-1} \text{ cm}^{-2}$ .



The colinear, dual, hydrogen-deuterium target enables us to collect high-precision, *in situ* calibration data so systematic uncertainties  $\leq 3\%$ .

Forward Detector	
Angular Range	$5^\circ - 40^\circ$ (outbenders)
$\Delta p/p$	$< 0.01$ @ 5 GeV/c
$\Delta\theta$ (mr)	$< 1$ $p > 2.5$ GeV/c
$\Delta\phi$ (mr)	$< 3$ $p > 2.5$ GeV/c
Neutron Detection	$N_{eff} = 0.1 - 0.6$

# More on the CLAS12 Detector

	Forward Detector	Central Detector
Angular Range		
Charged Particles	5° – 40°	40° – 135°
Photons	2° – 40°	N/A
Resolution		
$\Delta p/p$	< 0.01 @ 5 GeV/c	< 0.03 @ 0.5 GeV/c
$\Delta\theta$ (mr)	< 0.5	< 10
$\Delta\phi$ (mr)	< 0.5	< 6
Neutron Detection		
$N_{eff}$	0.1-0.6	0.1

# Beyond Elastic Form Factor Measurements

---

Additional form factor studies after the 12 GeV Upgrade.

Experiment	Spokesperson	Title	Hall	Beamtime
PR12-06-101	G. Huber	Measurement of the charged pion form factor to high $Q^2$	C	52 days
PR12-09-003	R. Gothe	Nucleon resonance studies with CLAS12	B	40 days


# Some More Background - Interpreting the EEFs

- At low momentum transfer ( $Q^2 \ll M_N^2$ )  $G_E$  and  $G_M$  are the Fourier transforms of the densities of charge and magnetization.


$$G_E(Q^2) = \int \rho(r) e^{-i\vec{q}\cdot\vec{r}} d^3r$$

where  $\vec{q}$  is the 3-momentum transferred by the electron.

- At high  $Q^2$  relativistic effects make the interpretation more interesting!



NSAC Long Range Plan



Arrington *et al.*, J.Phys.Conf.Ser. 299 (2011) 0120