

Physics 303

One-Dimensional Oscillators

1. An object of mass $m = 7.0 \text{ kg}$ is hung from the bottom end of a vertical spring fastened to an overhead beam. The object is set into vertical oscillations of period $T = 2.6 \text{ s}$. What is the force constant of the spring?
2. A particle of mass $m = 0.50 \text{ kg}$ is attached to a horizontal spring with a spring constant $k = 50 \text{ N/m}$. At the time $t = 0$, the particle has its maximum speed $v_{max} = 20 \text{ m/s}$ and is moving to the left. What is the particle's equation of motion? What is the minimum time interval required for the particle to move from $x = 0$ to $x = 1.0 \text{ m}$?
3. For the damped oscillator with $\gamma^2 < \omega_0^2$ show the general solution is

$$y(t) = c_1 e^{(-\gamma + i\Omega')t} + c_2 e^{(-\gamma - i\Omega')t} \quad (1)$$

where $\Omega' = \sqrt{\omega_0^2 - \gamma^2}$.

4. Apply the following boundary conditions

$$\text{for } t = 0 \implies y = y_0 \text{ and } \dot{y} = 0 \quad (2)$$

to Equation 1 and show

$$c_1 = y_0 \frac{\Omega' - i\gamma}{2\Omega'} \quad (3)$$

and

$$c_2 = y_0 \frac{\Omega' + i\gamma}{2\Omega'} \quad (4)$$

5. Now insert the results in Equations 3-4 into Equation 1 and show the following equation is true.

$$y(t) = \frac{y_0}{\Omega'} e^{-\gamma t} (\Omega' \cos \Omega' t + \gamma \sin \Omega' t) \quad (5)$$

6. Consider the function.

$$f(x) = \frac{1}{\sqrt{1+x}} \quad (6)$$

What is the Taylor series for this function for the first four terms? What does the n^{th} term look like? When can we approximate the function with the first two terms in the series? Explain.

7. For a damped oscillator, Newton's second law gives us

$$m \frac{d^2 x}{dt^2} = -kx - b \frac{dx}{dt} \quad (7)$$

in one dimension. Show that the expression

$$x = A e^{-bt/2m} \cos(\omega t + \phi) \quad (8)$$

is a solution as long as $b^2 < 4mk$.