Physics 303 One-Dimensional Oscillators

- 1. An object of mass $m = 7.0 \ kg$ is hung from the bottom end of a vertical spring fastened to an overhead beam. The object is set into vertical oscillations of period $T = 2.6 \ s$. What is the force constant of the spring?
- 2. A particle of mass $m = 0.50 \ kg$ is attached to a horizontal spring with a spring constant $k = 50 \ N/m$. At the time t = 0, the particle has its maximum speed $v_{max} = 20 \ m/s$ and is moving to the left. What is the particle's equation of motion? What is the minimum time interval required for the particle to move from x = 0 to $x = 1.0 \ m$?
- 3. For the damped oscillator with $\gamma^2 < \omega_o^2$ show the general solution is

$$y(t) = c_1 e^{(-\gamma + i\Omega')t} + c_2 e^{-(\gamma + i\Omega')t}$$

$$\tag{1}$$

where $\Omega' = \sqrt{\omega_0^2 - \gamma^2}$.

4. Apply the following boundary conditions

for
$$t = 0 \implies y = y_0$$
 and $\dot{y} = 0$ (2)

to Equation 1 and show

$$c_1 = y_0 \frac{\Omega' - i\gamma}{2\Omega'} \tag{3}$$

and

$$c_2 = y_0 \frac{\Omega' + i\gamma}{2\Omega'} \tag{4}$$

5. Now insert the results in Equations 3-4 into Equation 1 and show the following equation is true.

$$y(t) = \frac{y_0}{\Omega'} \ e^{-\gamma t} \left(\Omega' \cos \Omega' t + \gamma \sin \Omega' t\right) \tag{5}$$

6. For a damped oscillator, Newton's second law gives us

$$m\frac{d^2x}{dt^2} = -kx - b\frac{dx}{dt} \tag{6}$$

in one dimension. Show that the expression

$$x = Ae^{-bt/2m}\cos(\omega t + \phi) \tag{7}$$

is a solution as long as $b^2 < 4mk$.