



ELSEVIER

Contents lists available at ScienceDirect

## Environmental Research

journal homepage: [www.elsevier.com/locate/envres](http://www.elsevier.com/locate/envres)

# Prefrontal white matter pathology in air pollution exposed Mexico City young urbanites and their potential impact on neurovascular unit dysfunction and the development of Alzheimer's disease

Lilian Calderón-Garcidueñas<sup>a,b,\*</sup>, Rafael Reynoso-Robles<sup>c</sup>, Javier Vargas-Martínez<sup>b</sup>, Aline Gómez-Maqueo-Chew<sup>d</sup>, Beatriz Pérez-Guillé<sup>c</sup>, Partha S. Mukherjee<sup>e</sup>, Ricardo Torres-Jardón<sup>f</sup>, George Perry<sup>g</sup>, Angélica González-Maciél<sup>c</sup>

<sup>a</sup> The University of Montana, Missoula, MT 59812, USA

<sup>b</sup> Universidad del Valle de México, Mexico City 04850, México

<sup>c</sup> Instituto Nacional de Pediatría, Mexico City 04530 México

<sup>d</sup> Universidad del Valle de México, Hermosillo, Sonora 83299, México

<sup>e</sup> Mathematics Department, Boise State University, Boise, ID, USA

<sup>f</sup> Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, Mexico City 04310, México

<sup>g</sup> College of Sciences, University of Texas at San Antonio, San Antonio, TX, USA

## ARTICLE INFO

## Article history:

Received 1 October 2015

Received in revised form

30 November 2015

Accepted 27 December 2015

Available online 30 January 2016

## Keywords:

Air pollution

Alzheimer

BBB

Children

Dogs

Endothelial damage

PM<sub>2.5</sub>

Nanosize particles

Neurovascular unit

Tight junctions

## ABSTRACT

Millions of urban children are chronically exposed to high concentrations of air pollutants, i.e., fine particulate matter (PM<sub>2.5</sub>) and ozone, associated with increased risk for Alzheimer's disease. Compared with children living with clear air those in Mexico City (MC) exhibit systemic, brain and intrathecal inflammation, low CSF Aβ<sub>42</sub>, breakdown of the BBB, attention and short-term memory deficits, prefrontal white matter hyperintensities, damage to epithelial and endothelial barriers, tight junction and neural autoantibodies, and Alzheimer and Parkinson's hallmarks. The prefrontal white matter is a target of air pollution. We examined by light and electron microscopy the prefrontal white matter of MC dogs (*n*: 15, age 3.17 ± 0.74 years), children and teens (*n*: 34, age: 12.64 ± 4.2 years) versus controls. Major findings in MC residents included leaking capillaries and small arterioles with extravascular lipids and erythrocytes, lipofuscin in pericytes, smooth muscle and endothelial cells (EC), thickening of cerebrovascular basement membranes with small deposits of amyloid, patchy absence of the perivascular glial sheet, enlarged Virchow–Robin spaces and nanosize particles (20–48 nm) in EC, basement membranes, axons and dendrites. Tight junctions, a key component of the neurovascular unit (NVU) were abnormal in MC versus control dogs ( $\chi^2 < 0.0001$ ), and white matter perivascular damage was significantly worse in MC dogs (*p* = 0.002). The integrity of the NVU, an interactive network of vascular, glial and neuronal cells is compromised in MC young residents. Characterizing the early NVU damage and identifying biomarkers of neurovascular dysfunction may provide a fresh insight into Alzheimer pathogenesis and open opportunities for pediatric neuroprotection.

© 2015 Elsevier Inc. All rights reserved.

## 1. Introduction

Clinically healthy Mexico City (MC) children with no known risk factors for neurological or cognitive disorders exhibit cognition deficits, brain metabolic, structural and volumetric changes

\* Corresponding author at: The University of Montana, Missoula, MT 59812, USA.  
E-mail address: [lilian.calderon-garcidueñas@umontana.edu](mailto:lilian.calderon-garcidueñas@umontana.edu) (L. Calderón-Garcidueñas).

and the neuropathological and cerebrospinal fluid (CSF) laboratory hallmarks of Alzheimer and Parkinson's diseases i.e., tau hyperphosphorylation with pre-tangles, amyloid beta<sub>42</sub> (Aβ<sub>42</sub>) plaques, low CSF Aβ<sub>42</sub>, and misfolded α-synuclein accumulation (Calderón-Garcidueñas et al., 2008a, 2010, 2011a, 2012a, 2013a, 2015a).

Brain MRI and MRS studies in MC children and teens versus low air pollution controls show white matter metabolic changes and prefrontal white matter hyperintensities (WMH) (Wardlaw et al., 2013; Calderón-Garcidueñas et al., 2008b, 2011b, 2012b,

2015a, 2015b), while neuropathology findings reveal cortical disruption of the blood–brain barrier (BBB), endothelial activation, oxidative stress, high concentration of metals associated with combustion, inflammatory cell trafficking along with up-regulated gene network clusters including IL1, NFκB, TNF, IFN, and TLRs (Calderón-Garcidueñas et al., 2003, 2008a, 2009a, 2010, 2011a, 2013a, 2013b). Animal facility healthy MC young dogs also exhibit WMH by MRI, neuroinflammation, DNA oxidative damage, BBB breakdown, and accumulation of combustion-related metals (Calderón-Garcidueñas et al., 2002, 2003, 2008b, 2009a). The prefrontal cortex is a target of air pollution and its damage likely a major contributor to cognitive deficits in Mexico City young residents. Prior research has shown all epithelial and endothelial barriers are compromised in MC children and dogs and the production of high concentrations of endothelin-1 and autoantibodies against tight junction and neural proteins could be playing a role in the diffuse microvascular changes observed in young urbanites (Calderón-Garcidueñas et al., 2001, 2007, 2008c, 2009b, 2015d). In a recent study of 139 clinically healthy MC and control children age  $11.91 \pm 4.2$  y, serum antibodies against occludin/zonulin 1 and actin IgG, along with myelin oligodendrocyte glycoprotein, myelin basic protein, S-100, and cerebellar IgG were significantly higher in MC children (Calderón-Garcidueñas et al., 2015d). Zonula occludens (ZO) proteins are at the core of the protein networks which are anchored to the TJ-plaque dynamic structures and given that neuroinflammation is associated with BBB dysfunction and loss of tight junctions (Bauer et al., 2014; Elahy et al., 2015; Haseloff et al., 2015), we fully expected brain structural and metabolic changes in MC children.

Extensive data in the literature support human and animal breakdown of the nasal/olfactory, BBB and alveolar-capillary barriers and the expression of detrimental genes associated to urban air pollution (Harkema et al., 2006; Ljubimova et al., 2013; Van Miert et al., 2005; Kaplan et al., 2010; Carson et al., 2013; Bergin and Witzmann, 2013; Garwood et al., 2014). The work by Kamat et al. (2014), Winkler et al. (2014), Hawkes et al. (2014), Cabezas et al. (2014) and Garwood et al. (2014) is of particular interest to us given that their research support damage to brain endothelial cells occurs early in relation to Alzheimer's neuropathology and BBB disruption leads to neuronal damage, reactive gliosis, oxidative stress, neuroinflammation and early neurovascular dysfunction.

Of great concern in polluted environments with high concentrations of ultrafine particulate matter (UFP, nanosize particles < 100 nm) is that after passage through biological barriers, UFP end up in contact with the vascular endothelium and can induce damage (Wang et al., 2009; Gehr et al., 2011; Sharma et al., 2013; Ucciferri et al., 2014; Karmakar et al., 2014; Meng et al., 2015). The presence of high affinity autoantibodies against barrier forming proteins in urban children are critical to our understanding of air pollutant mechanistic damage pathways. There is robust evidence nanosize particles can increase endothelial paracellular permeability in vitro and induce endothelial TJ opening (Sharma et al., 2013; Yu et al., 2013; Ucciferri et al., 2014; Karmakar et al., 2014; Li et al., 2015).

The developing brain relies heavily on the delivery of oxygen and nutrients from the blood stream to meet metabolic demands of neural cells and blood supply, thus neural activity and vascular dynamics are tightly coupled (Lecrux and Hamel, 2011; Lacoste and Gu, 2015). The neurovascular unit (NVU) is the anatomical substrate of neurovascular interactions and a complex interaction between endothelial cells, pericytes, astrocytes, microglia and neurons is responsible for optimal delivery of oxygen and nutrients to the brain (Simard et al., 2003; Zlokovic, 2008; Lo and Rosenberg, 2009; Lacoste and Gu, 2015). A key function of this system is to keep a tightly control environment aimed to preserve the brain from toxins, pathogens and harmful chemicals.

Neurovascular dysfunction has a relevant focus in Alzheimer's disease (AD) research, particularly regarding BBB integrity, cerebral blood flow (CBF) and glucose transport into the brain (Iadecola, 2004; Keane and Campbell, 2015; Sweeney et al., 2015). White matter abnormalities are common in dementia and the pathology is the result of a combination of structural alterations of the cerebral vasculature, i.e., arteriosclerosis, cerebrovascular basement membrane pathology, and amyloid angiopathy, and nonstructural vascular abnormalities (vascular contractility or permeability) and/or neurovascular instability (Love and Miners, 2015).

Very little is known regarding the ultrastructural features of tight junctions (TJs), cerebrovascular basement membranes (Morris et al., 2014), capillaries, arterioles and axons in the prefrontal white matter of young healthy dogs and children with a lifetime exposure to urban air pollution. Our working hypothesis states that healthy, young dogs will have prefrontal vascular and white matter pathology and children living in the same area will share light and electron microscopic (EM) findings with those of dogs' raised in an animal facility in Mexico City.

We have one aim for this study: to document by 1 μm toluidine blue thick sections and EM the integrity of the prefrontal white matter in healthy young dogs resident in MC ( $n$ : 9) and in a cohort of MC children and teens ( $n$ : 26) autopsy prefrontal samples versus clean air controls (6 dogs and 8 children). Stored brain samples from seemingly healthy children dying suddenly in accidental deaths not involving the cranial cavity and undergoing forensic autopsies were the source of the frontal samples.

Our results identify abnormalities at the endothelial junctional complexes, microbleeds, perivascular lipid accumulation, abnormal cerebrovascular basement membranes, and the presence of ultrafine particles in mitochondria, basement membranes, axons and dendrites. Our study suggests that the integrity of the NVU in the prefrontal white matter is compromised in highly exposed young urbanites and short and long-term brain health consequences are expected.

## 2. Procedure

### 2.1. Study cities and air quality

Children's cohorts were selected from the Mexico City Metropolitan Area (MCMA) and control locations consisting of small cities in Mexico (Zacatlán and Huachinango, Puebla; Zitácuaro, Michoacán; Puerto Escondido, Oaxaca). The control cities have < 75,000 inhabitants and because of their small size their levels for the main criteria air pollutants (ozone, particulate matter, sulfur dioxide, nitrogen oxides and carbon monoxide) are lower than the current US EPA standards (Alonso et al., 2007).

Mexico City Metropolitan Area is an example of extreme urban growth and accompanying environmental pollution (Bravo-Alvarez and Torres-Jardón, 2002; Molina et al., 2010; Retama et al., 2015). The metropolitan area of over 2000 km<sup>2</sup> lies in an elevated basin 2200 m above sea level surrounded on three sides by mountain ridges. MCMA's nearly 24 million inhabitants, over 50,000 industries, and 5.5 million vehicles consume more than 50 million liters of petroleum fuels per day, producing an estimated annual emission of 2.3 million tons of particulate and gaseous air pollutants. MCMA motor vehicles produce abundant amounts of primary fine particulate matter (PM<sub>2.5</sub>). The high altitude and tropical climate where the MCMA is settled facilitate ozone production all year and contribute to the formation of PM<sub>2.5</sub>. Children from MCMA were residents in the northern-industrialized and southern-residential zones. Northern children have been exposed to higher concentrations of volatile and toxic organic compounds,

PM<sub>10</sub>, and PM<sub>2.5</sub> including high levels of its constituents: organic and elemental carbon, nitro- and polycyclic aromatic hydrocarbons and metals (Zn, Cu, Pb, Ti, Mn, Sn, V, and Ba), while southern children have been exposed continuously to significant and prolonged concentrations of ozone, secondary aerosols (NO<sub>3</sub>) and particulate matter associated with lipopolysaccharide PM-LPS.

## 2.2. Subjects

The work described in this research was conducted in accordance with the Code of Ethics involving animal research and the Institutional Animal Care and Use Committees. The frontal dog samples used in this work were obtained previously from two independent studies involving the use of Nimesulide<sup>®</sup> in mixed beagle dogs and a metal study (Calderón-Garcidueñas et al., 2002, 2003, 2008b, 2009a). Procedures used were in accordance with the guidelines of the Use and Care of Laboratory Animals (NIH Pub No. 86-23). The autopsy frontal samples were obtained from forensic cases with no identifiable personal data, not meeting the regulatory definition of human subject research.

### 2.2.1. Animal facility young dogs

Previously harvested dog frontal tissues for EM were used for this study. Two cohorts, one from MC and a control cohort of mixed beagles were whelped and housed in an outdoor–indoor kennel; husbandry was in compliance with the American Association of Laboratory Animal Certification Standards. Dogs were under daily veterinarian observation during their entire life, and at no time there was any evidence of respiratory, cardiovascular, gastrointestinal or neurological diseases. Dogs had all applicable vaccines and were treated with anti-helminthics regularly. Dogs from both cohorts had the same diets. Dogs were sacrificed with an overdose of sodium pentobarbital. Tissue blocks were archived in phosphate-buffered saline 0.1 M pH 7.5 with sodium azide at 4 °C prior to processing for EM. We selected to use frontal white matter tissue optimally fixed for electron microscopy from 15 dogs (3.17 ± 0.74 years). The 9 Mexico City selected dogs age 3.11 ± 0.67 years were in the non-treated Mexico City group exposed 24/7 to the Southwest MC atmosphere from birth. Six control dogs average age 3.23 ± 0.81 years were also studied (Table 1).

### 2.2.2. Light microscopy in dog samples

Frontal samples were post-fixed in 1% osmium tetroxide and embedded in Epon. Semi-thin sections (0.5–1 μm) were cut and stained with toluidine blue for light microscopic examination. Thirty blocks from the right (n: 15) and left (n: 15) prefrontal white matter were cut and examined in each dog. Each toluidine blue 1 μm section (30 slides in each dog) was examined under a microscope Carl Zeiss Axioskop 2 Plus equipped with a AxioVision REL 4.8 imaging system. For the measurement of abnormal neuropil areas we selected a square of 60 μm<sup>2</sup> and 16 areas of neuropil were reviewed and measured in 4 randomly selected toluidine blue 1 μm section slides (2R/2L) in each dog. One EM researcher was in charge of the measurement of the abnormal neuropil areas and the percentage of white matter damage was recorded in each dog, within each cohort, MC versus controls. This researcher in charge of the measurement of all slides was blind to the identification of the dog. A board-certified pathologist (LCG) without access to the identification codes reviewed the sections and the representative pictures.

### 2.2.3. Examination of dogs' frontal white matter samples by Transmission Electron Microscopy (TEM)

We selected the EM areas from the semi-thin sections stained with toluidine blue. Ultra-thin sections (60–90 nm) were cut and collected on slot grids previously covered with Formvar

**Table 1**

Summary of results comparing Mexico City with low air pollution control dogs<sup>a</sup>.

	Controls n:6	Mexico City n:9	p value
Age in years	3.23 ± 0.81 years	3.11 ± 0.67 years	0.95
Gender	3F/3M	5F/4M	NA
White matter percentage damaged area	0.77 ± 0.46	19.43 ± 2.69	0.002

<sup>a</sup> Measured in a 60 μm<sup>2</sup> area/16 areas from 4 slides 2R/2L.

membrane. Sections were stained with uranyl acetate and lead citrate, and examined with a JEM-1011 (Japan) microscope. Each electron micrograph was evaluated separately, then compared by group. We captured ultrastructural blood vessel images including sites of TJ's complexes, cerebrovascular basement membranes and the neuropil components. An average of 30 consecutive blood vessels and adjacent neuropil from each animal were documented. One EM researcher examined the TJs structures at magnifications between 12,000 × and 100,000 ×, took representative pictures and recorded normal and abnormal endothelial TJ structures from capillaries based on the following characteristics: observation of the endothelial junction undulated trajectory in a plane parallel to the basement membrane and exhibiting structurally closed and intact TJs as elegantly described by Muir and Peter (1962) and Castejón (2012, 2014), open inter-endothelial junctions, and loss of integrity and disruption of TJs.

We also noted astrocyte end feet detachment from the basement membranes, caveolar EC activity, the characteristics of the basement membranes, and microvascular endothelial cell damage including irregularity of the endothelial cell luminal surface, micro-villi like protusions, and bleb-like structures. The comparison between the results of the evaluation of 40 endothelial cell to endothelial cell TJs in each of the 15 animals, including 6 controls was assessed for significance with a chi square test.

### 2.2.4. Examination of children and teens prefrontal white matter samples by light and Transmission Electron Microscopy (TEM)

In the process of selecting the frontal samples we examined the entire autopsy in each subject and ruled out the presence of pathological processes other than the lesions associated to the sudden accidental death. The striking difference between dog and human frontal EM samples was the optimal preservation of tissues in dogs v humans given the time between death and autopsy. Given that the neuropil was not optimally preserved for EM in humans, the quantification of abnormal neuropil areas was not done. Frontal samples were post-fixed in 1% osmium tetroxide and embedded in Epon. Semi-thin sections (0.5–1 μm) were cut and stained with toluidine blue for light microscopic examination. Thirty blocks from the right (n: 15) and left (n: 15) prefrontal white matter were cut and examined in each child.

### 2.2.5. Statistical analysis

Statistical analyses were carried out using Excel and R (<https://www.r-project.org/>). All data are expressed as mean ± SD. We carried out statistical tests for intergroup differences after adjusting age. We used linear regression technique to accomplish this. Next, we tested for the significance of differences between the measurements of tight junction abnormalities in the control and Mexico City group by using the Chi-square test.


### 3. Results

#### 3.1. Air quality data


Mexico City residents are exposed year-round to particulate matter (PM) concentrations above United States National Air Quality Standards (NAAQS). For this work we focused on inhalable PM, broadly defined by the diameter of the aerodynamic particles, and classified into coarse particles ( $< 10$  to  $> 2.5$   $\mu\text{m}$ ;  $\text{PM}_{10}$ ), fine particles ( $< 2.5$   $\mu\text{m}$ ,  $\text{PM}_{2.5}$ ) and ultrafine PM (UFPM  $< 100$  nm). In spite of the efforts by the authorities to control PM air pollution, fine particles are still a health problem in MC because their levels have not shown a reduction trend in the last 10 years. Both, the  $\text{PM}_{2.5}$  annual air quality standard of  $12$   $\mu\text{g}/\text{m}^3$  and the 24-h standard of  $35$   $\mu\text{g}/\text{m}^3$  have been historically exceeded across the metropolitan area (Fig. 1). The highest concentrations occur in the NE sector (Xalostoc) where industrial and traffic activities are prevalent, and decrease towards the SW residential area (Pedregal).

As Fig. 2 shows, North-East MC teenagers older than 15 years old were conceived and born when  $\text{PM}_{2.5}$  average 24-hr levels were 2 to 3 times the current NAAQS. It should be noticed, every resident in MC has been exposed to  $\text{PM}_{2.5}$  concentrations above the respective NAAQS regardless of their place of residency within the city. Chemical PM composition studies in Mexico City have shown that the proportion of the different component PM species has not change significantly along the years (Bravo-Alvarez and Torres-Jardón, 2002; Vega et al., 2010; Molina et al., 2010; Retama et al., 2015). Fig. 3 shows the chemical composition of coarse ( $\text{PM}_{10}$ – $\text{PM}_{2.5}$ ) and fine ( $\text{PM}_{2.5}$  and  $\text{PM}_1$ ) particles in a representative site of MC.

In general,  $\text{PM}_{10}$  in MCMA is dominated by the fine fraction. The  $\text{PM}_{2.5}/\text{PM}_{10}$  ratio variations and the PM chemical composition are dependent on the site location and on the season. The  $\text{PM}_{2.5}/\text{PM}_{10}$  ratio shows relatively lower values at the E and NE sectors relative to the SW sites. Typically, the coarse PM is strongly dominated by geological material



**Fig. 1.** Box plot for 24-h  $\text{PM}_{2.5}$  concentrations at representative site of Mexico City Metropolitan Area: Pedregal (residential SW area) from 1997 to 2014. The dashed lines inside the box are the annual average and the continuous line the 24-h median. The  $\text{PM}_{2.5}$  annual mean NAAQS concentration value is represented by the dashed blue line and the  $\text{PM}_{2.5}$  24-h average NAAQS level is shown with the red continuous line.  $\text{PM}_{2.5}$  data were not available in MCMA until 2004, thus the  $\text{PM}_{2.5}$  trends were approximated using a correlation equation of the  $\text{PM}_{2.5}/\text{PM}_{10}$  ratio for the period 2004–2011 and  $\text{PM}_{10}$  data measured at each site of the period 1997–2003.  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  data were obtained from the Secretaría del Medio Ambiente del Distrito Federal (<http://www.aire.df.gob.mx>). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



**Fig. 2.** Box plot for 24-h  $\text{PM}_{2.5}$  concentrations at representative site of Mexico City Metropolitan Area: Xalostoc industrial and high traffic (NE area) from 1997 to 2014.

( $\text{SiO}_2 + \text{CO}_2^{-3} + \text{Al}_2\text{O}_3 + \text{Ca} + \text{Fe} + \text{Mg} + \text{K}$ ) from dust resuspension and its proportion diminishes as the particle decreases in size. In contrast, and of key importance for the brain effects, organic and carbonaceous aerosols are the dominant species in the PM fine fraction. Particle emissions from gasoline and Liquefied Petroleum Gas Combustion (LPG) are dominated by organic carbonaceous aerosols (OC), while in diesel particles black carbon (BC) is the key component. OC species include large alkanes, alkanolic acids, benzoic acids, benzaldehydes, phenols, alkanals, etc. (Seinfeld and Pandis, 1998). Contrary to expected, BC concentrations in  $\text{PM}_{2.5}$  have not shown a decrease through the years (Retama et al., 2015). BC is associated with polycyclic aromatic compounds (PAHs). PAHs are semivolatile species formed through the fusion of two or more benzene rings by a pyrolytic process during the incomplete combustion of carbonaceous fuels such as gasoline and diesel vehicle exhaust gases. Most of PAHs in MCMA are present in the fine fraction ( $\text{PM}_{2.5}$ ) contributing with 75–85% of the total mass. In general, low molecular PAHs have a higher ratio than high molecular PAHs in  $\text{PM}_{2.5}$  (Mugica et al., 2010). On the other hand, secondary inorganic aerosols typically comprise around 20% in all  $\text{PM}_{10}$  fractions. They are dominated by ammonium sulfate showing the highest levels at the Pedregal southwest area.

Trace metals in fine particles comprises around the 50% of the  $\text{PM}_{10}$ . The most abundant metals in  $\text{PM}_{2.5}$  are Zn, Cu, Pb, Ti, Sn, Ba, Mn, Sb, V, Se, As, Ni, Cd, and Cr in that order (Querol et al., 2008). Zn, Cu, Ba, Pb, Pb and Cd are tracers of road traffic, while V and Ni are tracers of industrial emissions. MC children in this study have been exposed to significant concentrations of  $\text{PM}_{2.5}$  during their entire life, including the prenatal period. The high concentrations of  $\text{PM}_{2.5}$  coincide with the time children play outdoors and/or stay in schools with broken windows and doors and are in close proximity to high traffic and fixed sources of pollution (Villarreal-Calderón et al., 2002). Children are also exposed to ozone concentrations above the USA standards (Calderón-Garcidueñas et al., 2015d). All other criteria pollutants for MCMA, including nitrogen dioxide, sulfur dioxide and lead have been at or below the current EPA standards (Secretaría del Medio Ambiente del Distrito Federal: <http://www.aire.df.gob.mx>). Control children have been lifelong residents in low pollution cities with all criteria air pollutants below the US EPA NAAQS standards (Calderón-Garcidueñas et al., 2012a).

#### 3.2. Quantification of the abnormal neuropil prefrontal areas in dogs'




Fig. 3. Average mass and composition of PM coarse (PM10–PM2.5), PM2.5 and PM1 at the Instituto Mexicano del Petróleo site (northern Mexico City) during March 2006 (based upon a compilation of Molina et al. (2010), Querol et al. (2008) and Aiken et al., 2009).

samples

Measurement of abnormal neuropil areas showed a significant difference between control and Mexico City dogs matched by age,  $p=0.002$  (Table 1).

3.3. Light microscopy and electron microscopic dogs' and children results

There was no statistical difference in the selected control versus MC dogs' ages ( $p=0.95$ ) Table 1 and control versus MC children ( $p=0.93$ ) (Table 2). Children carrying an APOE 4 allele were representative of the population at large (17.6%). The white matter pathology results in Mexico City dogs, children and teens versus controls were divided in 4 sections: (1) one micron toluidine blue light microscopy of blood vessels including capillary, postcapillary venules and arterioles (2) Endothelial ultrastructural pathology, including tight junctions (TJs), pericyte and smooth muscle cell morphology and cerebrovascular basement membranes, (3) Oligodendroglia, axonal, and dendritic electron microscopy and (4) Localization of nanosize particles.

3.3.1. Light microscopy morphological assessment using toluidine blue staining in dog's frontal white matter

Control samples of frontal white matter exhibited normal blood vessels ranging from arterioles, capillaries and venules. Virchow-Robin (VR) spaces were unremarkable and free of mononuclear cells, red blood cells (RBC) or lipid droplets. The neuropil was unremarkable and in particular the myelinated axons varied in size, they were uniformly distributed and intact (Fig. 4A). In contrast, Mexico City dogs exhibit expanded VR spaces with mononuclear cells characterized by large nuclei with increased chromatin condensation (Fig. 4B). We observed patchy pallor of the neuropil with zones where the myelinated axons were scanty and alternated with clusters of small myelinated fibers (Fig. 4C), along extensive areas of rarefaction of the neuropil around blood vessels (Fig. 4D\*). Large perivascular cells with the morphology of macrophages and clusters of lipid droplets were prominent around blood vessels (Fig. 4E and F). Distributed throughout the frontal white matter, small blood vessels exhibit hyperplastic endothelial cells reducing their lumen (Fig. 4G). In the transition between the subcortical white and the gray matter, abnormal small vessels were also present with RBC occupying the perivascular space (Fig. 4H). Remarkably, neuronal bodies in the transition area exhibited vacuolated cytoplasm.

3.3.2. Light microscopy morphological assessment using toluidine blue staining in children's frontal white matter

Control children exhibited unremarkable blood vessels,

Table 2


Summary of controls and Mexico City children and teens selected for light and electron microscopy prefrontal samples.

Controls age	Controls gender	Controls Apoe
8.2	1	34
17	1	33
13	1	33
7.8	0	33
16	1	34
14	1	33
15	1	33
10	0	33
12.62 ± 3.5 years	6M/2F	2 Apoe 3/4
MC age	MC gender	MC apoe
15	1	44
13	0	34
15	1	44
18	1	44
17	1	33
11	1	33
16	1	33
2	1	33
17	1	33
17	1	33
12	1	33
3	1	33
17	1	33
17	1	33
11	0	33
14	1	33
14	0	33
1	1	33
7	0	33
13	1	33
15	1	33
14	0	33
13	0	33
11	0	33
15	0	33
16	1	33
12.67 ± 4.9 years	18M/8F	4 Apoe4

Table 3

Summary of TJs observations in endothelial capillary prefrontal dogs' vessels. Results yielded a  $\chi^2$  statistic=21.96, degrees of freedom=1,  $\chi^2 < 0.0001$ .

Control dogs n:6	Mexico city dogs n:9
40 TJs observations in each dog	40 TJs observations in each dog
240 TJ	360 TJ
225 normal	288 normal
15 abnormal	72 abnormal




**Fig. 4.** Light microscopy assessment using toluidine blue staining in dog's frontal white matter. Unremarkable white matter is characteristic of control dogs (A). Virchow–Robin (VR) spaces are free of mononuclear cells, red blood cells or lipid droplets. The neuropil is unremarkable and the myelinated axons (A) vary in size, they are uniformly distributed and they are intact. In contrast, Mexico City dogs exhibit expanded VR spaces with mononuclear cells characterized by large nuclei with increased chromatin condensation (B, arrowheads). Scattered through the neuropil, cells with large nuclei and increased chromatin condensation are present (B short arrows). Patchy pallor of the neuropil with zones where the myelinated axons are scanty and alternate with clusters of small myelinated fibers (C square) and cells showing prominent chromatin condensation (C arrowheads) are common in MC dogs. Small arterioles show thick walls and expanded VR spaces (long arrows). In D we observed extensive areas of rarefaction of the neuropil around blood vessels (\*). Large perivascular cells have the morphology of macrophages (arrowheads), while clusters of lipid droplets are prominent around blood vessels (long arrows). In E, a close up of a typical blood vessel in a young dog showing large accumulation of lipids (arrowheads). F shows a postcapillary venule with perivascular lipid accumulation (arrowheads). Distributed throughout the frontal white matter, small blood vessels (G) exhibit hyperplastic endothelial cells (arrowheads) reducing their lumen (L). Mononuclear cells are seen attached to the hyperplastic endothelium (long arrow). In the transition between the subcortical white and gray matter (H), abnormal small vessels are also present with RBC occupying the perivascular space (arrowheads). Remarkably, neuronal bodies in the transition area have vacuolated cytoplasm (short arrows).

Virchow–Robin (VR) spaces and neuropil (Fig. 5A). MC children showed significant expansion of the VR spaces and clusters of perivascular lipids (Fig. 5B and C). Scattered small tortuous blood vessels showed mononuclear cells in the expanded perivascular spaces and leaking of lipid material affected small capillaries (Fig. 5D and E). A few of the blood vessels architecture was completely lost (Fig. 5F). Axonal abnormalities were present and a few large axons show intraaxonal vacuolation and curled membrane fragments. Hyperplastic endothelial cells were common (Fig. 5G). In the transition between the subcortical white and the gray


matter, abnormal small vessels were present with amorphous material occupying the intra and perivascular spaces (Fig. 5H).

### 3.3.3. Electron microscopy of dogs' blood vessels and neuropil

Mexico City dogs exhibited abnormal small blood vessels with irregular endothelial basement membranes, RBC in the widened perivascular spaces along with clusters of lipid material and cell fragments (Fig. 6A and B). Early formation of lipofuscin was a common finding in pericytes of < 2y old dogs, while capillaries showed irregular luminal endothelial cell surfaces with microvilli-



**Fig. 5.** Light microscopy assessment using toluidine blue staining in children's frontal white matter. Control children (A) exhibit unremarkable blood vessels, Virchow–Robin (VR) spaces and neuropil. MC children show lipid material accumulation around blood vessels (B\*) and a significant expansion of the VR space and clusters of perivascular lipids (C, short arrows). Scattered small tortuous blood vessels show mononuclear cells (D, arrowheads) in the expanded perivascular spaces (\*). A few large axons (A) show intraaxonal vacuolation (short arrows) and curled membrane fragments (long arrows). Leaking of lipid material also affects small capillaries (E\*). A few of the blood vessels architecture is completely lost (F). Scattered unidentified cells show clumping of the chromatin (short arrows). Hyperplastic endothelial cells are also seen (G, arrowheads). In the transition between the subcortical white and gray matter, abnormal small vessels are present with amorphous material occupying the intra and perivascular spaces (H\*).



**Fig. 6.** Electron microscopy of dogs' blood vessels. An abnormal small blood vessel (A) with irregular endothelial basement membranes (arrowheads) and an apparently intact perivascular glial sheath (short arrows) shows clusters of lipid material (\*). RBC are seen in the perivascular space. B shows a capillary vessel with a wide perivascular space (\*) occupied by cell fragments (short arrows). Note the significant rarefaction of the neuropil on the right side of the picture (square). Early formation of lipofuscin is seen in pericytes of < 3y old dogs (C short arrow). Capillaries show irregular luminal endothelial cell (EC) surface with microvilli-like protrusions (long arrow) and a large perivascular space with abundant cellular debris (\*). A close up of the lipofuscin formation can be seen in D. A lipofuscin granule (Lf) with dense osmiophilic content, and one mitochondria with abnormal cristae (D, short arrow) lie in the adjacent cytoplasm. In E, a small vessel shows a thick and irregular basement membrane (arrowheads). Scattered capillaries show cells in the position of pericytes and smooth muscle cells with semicircular or circular thickening of vascular walls containing large amounts of amorphous material (F\*) and smaller areas with amyloid-like fibrils (short arrows). The endothelial cell shows microvilli-like protrusions (long arrow) and there is a white blood cell (WBC) and one RBC occupying the vessel lumen (L). G shows helical twist amyloid fibers with a 35–50 Å repeat (short arrow). A blood vessel with a prominent endothelial cell nucleus (EC) protruding in the lumen is seen in H along with RBCs. A smooth muscle cell (SMC) cytoplasm display multiple mitochondria (short arrows). An unidentified perivascular structure shows a vacuolated background (\*) with cellular fragments (short arrows). A capillary with an attached white blood cell (WBC) occupies the lumen of the vessel (I).


like protrusions (Fig. 6C and D). Thick and irregular basement membranes were a common finding (Fig. 6E). Scattered capillaries showed cells in the position of pericytes and smooth muscle cells with semicircular or circular thickening of vascular walls containing large amounts of amorphous material and smaller areas with amyloid-like fibrils (Fig. 6F and G). Endothelial cells showed microvilli-like protrusions, prominent endothelial cell nuclei protruding in the lumen and attached white blood cells (WBC) occupying the lumen of the vessels (Figs. 6H and I).

White matter arterioles displayed layers of smooth muscle cells (Fig. 7A and B). Tight junctions between endothelial cells were mostly unremarkable in Mexico City dogs (Fig. 7C). A significant number of prefrontal white matter capillaries in highly exposed dogs were characterized by wide VR spaces and the patchy absence of astrocytic perivascular endfeet (Fig. 7D). Mononuclear cells commonly occupied the VR space and projected cytoplasmic

processes towards the capillary wall while patchy areas of neuropil were characterized by the absence of axons and cellular profiles. Unremarkable tight junctions alternated with the presence of osmiophilic granular material obliterating the cleft of the TJs and focal lack of integrity of the TJs (Fig. 7F and G). In contrast, control animals displayed unremarkable TJs (Fig. 7H). Quantification of the abnormal tight junction in prefrontal endothelial cells yielded a significant difference between control and Mexico City dogs,  $\chi^2$  statistic = 21.96, degrees of freedom = 1,  $\chi^2 < 0.0001$  (Table 3).

### 3.3.4. Electron microscopy of children' blood vessels and neuropil

The typical prefrontal findings in Mexico City children included: small blood vessels with accumulation of lipofuscin in pericytes and undulating irregularly thickened EC basement membranes (Fig. 8A, B, and C). Striking isolated arteriolar white matter changes were observed in Mexico City children (Fig. 8D).



**Fig. 7.** Electron microscopy of dogs' blood vessels. White matter arterioles display layers of smooth muscle cells (SMC) (A). Endothelial cells (short arrows) and RBC are seen. In a close-up (B), the typical components of a smooth cell cytoplasm are present, bundles of microfilaments (short arrows), elastic microfibrils (arrowheads), dense bodies and mitochondria (long arrows). A close-up of a tight junction between two endothelial cells (EC) is seen in C (arrowheads). A significant number of prefrontal white matter capillaries (D) in MC dogs are characterized by wide VR spaces and the patchy absence of astrocytic perivascular endfeet (arrowheads). Mononuclear cells (short arrows) occupy the VR space and project cytoplasmic processes towards the capillary wall (long arrows). Patchy areas of neuropil (\*) are characterized by the absence of axons and cellular profiles. E is a higher power of the square in D to focus on tight junctions in endothelial cells. A series of tight junctions (short arrows) between endothelial cells define the limits between cells. A RBC is seen in the lumen of the capillary. A higher power of the TJs (F) illustrates the clear lining of the cleft (white arrowheads) alternating with the presence of osmiophilic granular material obliterating the cleft of the TJs (short arrows). A portion of the RBC is seen in the lumen of the capillary. Higher magnification (150,000 $\times$ ) shows a TJs intact segment (G arrowhead) and the lack of integrity of the TJs (short arrows). In H a TJs (arrowheads) is intact in a control animal (RBC in lumen).

Changes included irregular and thick basement membranes and endothelial cells with large lysosomal bodies (Fig. 8E). Interspersed scant pyknotic nuclei were identified in relation with smooth muscle cells. Hyperplastic endothelium associated with a reduction of the lumen, marked thickening of the endothelial basement membranes and extensive perivascular areas of cell debris, vacuolization of the neuropil and abnormal large axons were common findings in MC samples, but not in controls (Fig. 8F). Irregular basement membrane varied in thickness between 0.8 and 2  $\mu\text{m}$  (Fig. 8G and H). Occasionally, we observed the apparent penetration of a mononuclear luminal cell through the endothelium (Fig. 8I and J) and a poor focal definition of the BM endothelial cell (Fig. 8K).

Nanosize particles could be seen in the endothelial cell cytoplasm close to the basement membranes and abundant caveolar activity was present in endothelial cells of MC children (Fig. 8L) and strikingly absent in controls. Nanosize PM was seen in RBC, endothelial cell mitochondria, basement membranes and abnormal mitochondria (Fig. 9A–E). Common observations between RBC and the endothelial cell cytoplasm was the presence of linear accumulation of nanosize particles (Fig. 9G and H).

#### 4. Discussion


Vascular and perivascular damage in the prefrontal white matter is a major feature of young Mexico City residents exposed to concentrations of fine particulate matter and ozone above the current USA standards. Major findings including abnormal TJs in endothelial cells, leaking of capillaries, abnormal cerebrovascular

basement membranes, and patchy absence of the perivascular glial sheet are likely compromising the integrity of the neurovascular unit. Tight junctions, a critical component of the NVU were abnormal in MC compared to control dogs, and white matter perivascular damage was significantly worse in MC *healthy* dogs.

The potential consequences of this evolving neurovascular unit pathology in urban children and dogs are highly relevant and pivotal to the pathogenesis of vascular-based neurodegenerative disorders like Alzheimer's disease (Zlokovic, 2008; Lo and Rosenberg, 2009; Wardlaw et al., 2013; Winkler et al., 2014; Garwood et al., 2014; Weinl et al., 2015; Elahy et al., 2015; Qosa et al., 2015; Keaney and Campbell, 2015; Tietz and Engelhardt, 2015).

Dysfunction of the NVU in childhood can ultimately lead to devastating short and long term consequences including dysregulation of cerebral blood flow, focal vascular insufficiency, innate immunity dysregulation, neuroinflammation, microstructural altered properties of major fiber tracts and regional volumes of white matter, alterations in ABC efflux transporters expression and/or activity, failure of elimination of interstitial fluid from white matter, vascular deposition of amyloid- $\beta$  and failure in the flow along pericapillary basement membranes to supply nutrients to the neuropil and equally important, to drain out waste products and soluble metabolites (Weller et al., 2009, 2015; Lo and Rosenberg, 2009; Mathiisen et al., 2010; Hawkes et al., 2014; Keaney and Campbell, 2015; Qosa et al., 2015; Tietz and Engelhardt, 2015; Brickman et al., 2015; Sweeney et al., 2015). The crosstalk between TJ and adherens junctions to maintain barrier integrity is likely to be disrupted, altering CNS homeostasis (Tietz and Engelhardt, 2015; Weinl et al., 2015; Keaney and Campbell, 2015).


There are numerous studies recognizing cerebrovascular



**Fig. 8.** Electron microscopy of children's prefrontal white matter. A prefrontal white matter small blood vessel (A) exhibits a typical finding in MC children: accumulation of lipofuscin (arrowheads) in pericytes. Extensive accumulation of lipofuscin (*Lf*) is a major finding in perivascular cells (B). A pericyte is present (arrowheads), along with a RBC occupying the entire vessel lumen. In C, an oligodendroglial cell (OLIGO) is seen adjacent to a small blood vessel with an undulating basement membrane (arrowheads). Striking arteriolar white matter changes are observed in MC teens (D). An arteriole exhibits an irregular and thick basement membrane (short arrows) between the endothelial cells (EC) and the tunica media layer with numerous smooth muscle cells (SMC). A higher power (E) shows EC with large lysosomal bodies (arrowheads), irregular and thick basement membrane (short arrows) and the concentric rings of smooth muscle cells (SMC). Interspersed scant pyknotic nuclei are identified in relation with smooth muscle cells (long arrows). A smaller arteriole (F) shows a hyperplastic endothelium (EC) with a reduction of the lumen (L), marked thickening of the endothelial basement membranes (arrowheads) and extensive perivascular areas of cell debris, vacuolization of the neuropil (VAC) and abnormal large axons (A). In G, EC's are resting on an irregular basement membrane (arrowheads). Variations in the thickness of the basement membrane are significant (H, arrowheads). In the same vessel, another feature includes the apparent penetration of a mononuclear luminal cell (MC) through the endothelium (I). The nucleus elongates and indents the basement membrane (short arrow). Notice the close apposition of the mononuclear and the endothelial cell (I, long arrows and J). Four punctate junction structures are seen between the endothelial and the mononuclear cell (short arrows). A close-up of the basement membrane on the abluminal side of the EC shows a poor definition of the BM limits (K arrowheads). The endothelial cell shows a large lysosomal body (\*). In L a 22 nm particle (short arrow) is seen in the endothelial cell cytoplasm close to the basement membrane. There are abundant pinocytotic vesicles in the EC cytoplasm (long arrows).

contributions to neurodegenerative diseases with both neurovascular and neurometabolic uncoupling playing a role in age-related neuropathologies in close association with cognitive decline (Brickman et al., 2015; Lourenco et al., 2015; Haight et al., 2015; Lai et al., 2015). The statement by Lo and Rosenberg (2009) that *sick vessels may generally lead to sick neurons* is of great concern in a developing brain. In the complex scenario of lifetime multiple air pollutant exposures, both the ones we usually measure and the ones we do not, we have plenty of evidence that particulate matter in the nanosize range can induce endothelial damage (Ucciferri et al., 2014; Alinovi et al., 2015; Bayat et al., 2015; Spigoni et al., 2015; Meng et al., 2015). Nanoparticle exposure is associated to systemic immune responses (Thompson et al., 2014) and endothelial heterogeneity in response to exogenous stimuli, particle potency and composition, sources and the particle content of

metals and endotoxin are key in the evaluation of endothelial damage and inflammatory responses and the severity of oxidative stress (Karthikeyan et al., 2013; Alinovi et al., 2015). Metals are a key component of particles and Mexico City children have high concentrations in the frontal cortex of manganese, nickel and chromium parallel to high expression of COX2, IL1 $\beta$  and TGF- $\beta$ 1 indicating neuroinflammation (Calderón-Garcidueñas et al., 2012a, 2013a, 2013b). IL1 $\beta$ , a major microglial pro-inflammatory cytokine has proven to be an early marker of air pollution exposure in MC children's brains (Calderón-Garcidueñas et al., 2008a), and its role in BBB disruption through downregulation of Sonic hedgehog (SHH) in astrocytes along increased production of pro-inflammatory chemokines including CCL2, CCL20 and CXCL2 are critical for BBB disruption and neuroinflammation (Wang et al., 2014). Up-regulation of TGF- $\beta$ 1 is a common finding in frontal



**Fig. 9.** Localization of nanosize particles in the white matter. A illustrates nanosize particles in the RBC, an endothelial cell mitochondria (M) and the basement membrane (arrowheads). Particles in the range of 20–38 nm are seen in abnormal mitochondria (M) with a few cristae (arrowheads) or no cristae (\*) (B). An intact mitochondria displaying a couple of particles (short arrows) is seen within a myelinated axon (C). In contrast, a poorly preserved unmyelinated axon (arrowheads) shows a larger particle measuring 40 nm (long arrow). In an adjacent area (D), a poorly preserved unmyelinated axon (arrowheads) shows a mitochondria with no particles. Poorly preserved unmyelinated axons show mitochondria with intact cristae and one single particle 30 nm in diameter (E, short arrow). A degenerating myelinated axon with remnants of myelin (arrowheads) shows one single nanosize particle (short arrow) (F). A common observation between RBC and the endothelial cell cytoplasm is the presence of linear accumulation of nanosize particles (short arrows) (G and H). In H, a common finding in MC subjects: significant caveolar EC activity (long arrows).

cortex of MC children (Calderón-Garcidueñas et al., 2012a) and chronic production of TGF- $\beta$ 1 in transgenic mice has been associated with AD-like microvascular degeneration (Wyss-Coray et al., 2000).

Both endothelial cell injury and dysfunction are considered early events in air pollution-associated effects and the major marker of endothelial dysfunction: endothelin-1 is quickly produced in response to the combination of particulate matter and ozone (Thomson et al., 2005; Calderón-Garcidueñas et al., 2007; 2008c; Finch and Conklin, 2015). Endothelin-1 is a 21 amino acid peptide with potent vasoconstrictor properties. Vasoconstriction is a potential contributor to cerebral hemodynamic impairment in urban children that can be included in the context of systemic inflammation, vasospasm and induction of focal ischemia (Eisenhut, 2014; Murray et al., 2014; Lei et al., 2015; Nguemeni et al., 2015). ET-1 overexpressing astrocytic cells showed amyloid secretion after a hypoxic/ischemia insult, suggesting astrocytic ET-1 contributes to dementia associated with ischemic stroke (Hung et al., 2015).

Since the major factor determining the impact of neural antibodies is the integrity of the blood–brain-barrier (Diamond et al., 2014; Levin et al., 2010), the autoimmune TJs and neural components seen in MC children potentially add to the ongoing neurovascular unit damage. Complicating the white matter damage scenario is the issue of impaired drainage of interstitial fluid (Laman and Weller, 2013; Weller et al., 2015). MC children have white matter abnormalities on magnetic resonance imaging (MRI) including white matter hyperintensities (WMH) and visible perivascular spaces (PVS) (Calderón-Garcidueñas et al., 2011b, 2012b) and although there is very effective lymphatic drainage for fluid and solutes along restricted pathways in the basement membranes of cerebral capillaries and arteries in young individuals we

fully expect lymphatic drainage of the brain is impaired in MC subjects particularly in Apolipoprotein  $\epsilon$ 4 carriers (Weller et al., 2015). Weller et al., strongly suggest that deposition of proteins in the lymphatic drainage pathways in the walls of cerebral arteries relates to vascular pathology, including cerebral amyloid angiopathy and protein elimination failure angiopathy. In addition, the presence of abnormal proteins, including A $\beta$  in the walls of capillaries and arteries in Mexico City teens and young adults (Calderón-Garcidueñas et al., 2012a, 2013a) may suggest that A $\beta$  is deposited in the pathways by which interstitial fluid drains from the brain as described with great detail by Hawkes et al. (2014). The contribution of the neuropil pathology including the perivascular axonal and dendritic findings cannot be ignored given the strong evidence of the key role of cerebral small vessel vasculopathy to the pathogenesis of Alzheimer's disease (Edwards et al., 2015) and the work of Maillard et al., (2011) showing that WMHs and their penumbra vary in severity and together span a continuous spectrum of white matter injury that worsens with time. Promjunyakul et al., (2015) measured cerebral blood flow (CBF) surrounding WMHs to identify 'WM at risk'. They concluded CBF penumbra involves surrounding WMHs, which is associated with future WMH expansion. Prefrontal WMH's are seen in approximately 50% of seemingly healthy Mexico City children and young dogs (Calderón-Garcidueñas et al., 2008b), so the issue is a serious one, potentially affecting around 4 million children in MC alone.

The unfortunate combination of white matter hyperintensities, amyloid  $\beta$  and factors playing a role in modifying cerebral autoregulation in Mexico City children account for vascular risk factors for the development of AD (Calderón-Garcidueñas et al., 2012b, 2013a, 2015a). The white matter damage we are describing in this work could be directly related to two important findings in Mexico City children: (1) The presence of predominantly frontal white

matter hyperintensities (WMH) in MC children and their relationship with the complex modulation of cytokines and chemokines and cognitive correlates (Calderón-Garcidueñas et al., 2012b). (2) The statistically significant reduction of the NAA/Cr ratio in the right frontal white matter and decrements on attention, short-term memory, and below-average scores in Verbal and Full Scale IQ ( $> 10$  points) in APOE 4 v APOE 3 children (Calderón-Garcidueñas et al., 2015b). Interestingly, young dogs with frontal WMH expand their lesions with time in a strong association with arteriolar pathology (Calderón-Garcidueñas et al., 2008b). The issue of WMH in MC children is very important because white matter microstructure alterations in childhood will result in changes in fluid intelligence later in life (Ritchie et al., 2015). In fact, we are seeing fluid intelligence/working memory changes in elementary school children, teens and in young college students (Calderón-Garcidueñas et al., 2008b, 2015b and unpublished work on progress with Dr. Randall Engle at Georgia Tech). Moreover, WMH relate to cognitive and functional impairment (Prins and Scheltens, 2015).

At the core of the matter is the complex interaction between abnormal tight junctions, BBB dysfunction, small vessel cerebrovascular disease, abnormal cerebrovascular basement membranes, altered cerebral autoregulation, high production of ET-1, nanoparticles damaging endothelial cells, autoimmunity to tight junction and neural proteins, neuroinflammation and lifestyle and vascular risk factors at play in Mexican youth. Of particular concern is childhood obesity and overweight problems with poor diets and high consumption of high fructose sodas. We are observing 11 year old normal weight MC children with 12 h fasting hyperleptinemia related to PM<sub>2.5</sub> exposures, altered appetite-regulating peptides, and increases in ET-1 (Calderón-Garcidueñas et al., 2015e). These changes could signal the future trajectory of urban children towards the development of insulin resistance, obesity, type II diabetes, addiction-like behavior, cognitive impairment and Alzheimer's disease. Maternal obesity is also in the picture, since the risk for AD will be associated with sustained high-fat diet during gestation and lactation periods, as shown in a mouse model (Hawkes, 2015). Using a mouse model of maternal obesity, Hawkes et al. found that exposure to a high-fat diet in the fetus and the early neonatal period induced changes in multiple components of the neurovascular unit, including a down-regulation in collagen IV, fibronectin and apolipoprotein E, an up-regulation in markers of astrocytes and perivascular macrophages and altered blood vessel morphology in the brains of adult mice. Interestingly, sustained high-fat diet over the entire lifespan resulted in additional decreases in levels of pericytes and impaired perivascular clearance of A $\beta$  from the brain. These results are very relevant to human populations and support a critical role for early dietary influence on the brain vasculature across the lifespan, with consequences for the developing brain and later cerebrovascular and neurodegenerative diseases (Hawkes, 2015).

Two other potential consequences of the NVU pathology are worth commenting in the scenario of children living in polluted environments. The first one is nicely discussed by Licht and Keshet (2015) and pertains to the key issue relating blood vessels to stem cell niches in the brain. The sub-ventricular zone and the hippocampus are the two major neurogenic niches and stem cells are physically associated with blood vessels. Thus, our expectations will include an alteration of the NVU components in these two target areas in highly exposed children. Parenthetically, MC children of the same age as the children in this study show significant low NAA/Cr hippocampal ratios, but not yet robust markers of hippocampal inflammation (IL1 $\beta$  and COX2) (Calderón-Garcidueñas et al., 2008a, 2015c). The second issue has been discussed by several authors including Sweeney et al., and Poggessi et al., 2015 and deals with a pressing need: if indeed the NVU pathology is one

key factor in the development of AD then early identification of NVU cell-specific responses including endothelial dysfunction circulating biomarkers ought to be explored in highly exposed children. In this regard, the use of non-invasive techniques to study the NVU integrity are a must, since we are dealing with clinically healthy children. Arterial spin labeling (ASL) a non-invasive MRI technique to measure cerebral blood flow (CBF) could be a useful tool (Grade et al., 2015), while diffusion tensor imaging focusing on diffusivity measures could be used to assess white matter integrity (Nir et al., 2013) along with characterizing major global WM network properties (Fischer et al., 2015). Cerebrospinal fluid markers of NVU dysfunction are critical to study in exposed paediatric populations and a useful source of CSF samples are those labeled as normal in the course of exploratory studies, i.e., work-up for CNS leukaemia involvement (Calderón-Garcidueñas et al., 2015a). Sweeney et al. (2015) have a table depicting CSF markers of BBB breakdown, vascular cells and astrocytes that can be applicable to children's samples from low versus highly polluted places.

#### 4.1. Limitations, perspectives, and future directions

We acknowledge our main limitations include the relatively small group of dogs with optimal EM samples, the lack of optimal preservation of human frontal tissues due to the time between the death of the subjects and the processing of samples, the need to quantitate at a molecular level the mRNA and protein levels of the integral TJs proteins and the flow cytometry analysis of endothelial TJs proteins, inflammatory cytokines and adhesion molecules (Elahy et al., 2015).

Taken together, findings summarized in this work lend strong support to the notion that key components of the neurovascular unit are compromised in highly exposed Mexico City children and young dogs. Advancing from the state of 'guilt by association' in terms of air pollution components and their specific role on the brain impact we are describing, to identification of specific factors altering the NVU is exposed children are critical research areas. Preceding this paramount undertaking, however, further characterization of the biomarkers of neurovascular dysfunction might prove instrumental. Future research needs to address fundamental questions: how early are the white matter changes in the overall MC pediatric population? Are we able to pick up the differences between MC and controls in terms of DTI diffusivity? How early? Are APOE4 carriers different from APOE 3 from early childhood in terms of NVU damage markers? What is the impact of the lifestyle factors in the developing of NVU dysfunction? Are overweight and obese children at higher risk? Or perhaps normal weight children with hyperleptinemia are at high risk as well?

Independently of issues concerning the impact of air pollutants and what specific air pollutants are responsible within the complexity of the atmospheric chemistry and the microenvironments of each subject, there is a pronounced need to identify molecular, structural and functional phenotypes of children and teens associated with cognition deficits and neurodegenerative features because intervention to protect them is a must. Reliable neurovascular dysfunction biomarkers are essential for early detection of subjects at risk and intervention and to evaluate the effectiveness of early neuroprotection (Sweeney et al., 2015).

In the broad sense, current knowledge should allow us the opportunity to intervene in young urban populations, especially because we have enough evidence that multidimensional interventions are useful and because we cannot ignore the 138% increased risk of AD per increase of 4.34  $\mu\text{g}/\text{m}^3$  in PM<sub>2.5</sub> in the Jung et al. (2015) work suggesting long-term exposure to O<sub>3</sub> and PM<sub>2.5</sub> is associated with increased the AD risk.

Everybody talks about the enormous health, family, social, and

economic burden of Alzheimer's disease, why then ignore the fact that air pollution is likely playing a key role, along with lifestyle and genetic factors and that children's detrimental brain effects have potential serious consequences in the short and long term. There is no support for preventive research. Why?

### Financial support

None.

### Conflict of interest

None.

### References

- Aiken, A.C., Salcedo, D., Cubison, M.J., Huffman, J.A., DeCarlo, P.F., Ulbrich, I.M., Docherty, K.S., Sueper, D., Kimmel, J.R., Worsnop, D.R., Trimborn, A., Northway, M., Stone, E.A., Schauer, J.J., Volkamer, R.M., Fortner, E., de Foy, B., Wang, J., Laskin, A., Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N.A., Paredes-Miranda, G., Arnott, W.P., Molina, L.T., Sosa, G., Jimenez, J.L., 2009. Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (T0) – Part 1: fine particle composition and organic source apportionment. *Atmos. Chem. Phys.* 9, 6633–6653.
- Alinovi, R., Goldoni, M., Pinelli, S., Campanini, M., Aliatis, I., Bersani, D., Lottici, P.P., Iavicoli, S., Petyx, M., Mozzoni, P., Mutti, A., 2015. Oxidative and pro-inflammatory effects of cobalt and titanium oxide nanoparticles on aortic and venous endothelial cells. *Toxicol. Vitro* 29, 426–437. <http://dx.doi.org/10.1016/j.tiv.2014.12.007>.
- Alonso, E., Martínez, W., Rubio, J.C., Velasco, F., Chávez, H.L., Ávalos, M., Lara, C., Cervantes, E., 2007. Calidad del aire en cuatro ciudades de Michoacán, México: Su efecto sobre materiales de construcción. *Rev. De La Constr.* 6, 66–74.
- Bauer, H.C., Krizbai, I.A., Bauer, H., Traweger, A., 2014. You shall not pass" – tight junctions of the blood brain barrier. *Front. Neurosci.* 8, 392. <http://dx.doi.org/10.3389/fnins.2014.00392>.
- Bayat, N., Lopes, V.R., Schölermann, J., Jensen, L.D., Cristobal, S., 2015. Vascular toxicity of ultra-small TiO<sub>2</sub> nanoparticles and single walled carbon nanotubes and in vivo. *Biomaterials* 63, 1–13. <http://dx.doi.org/10.1016/j.biomaterials.2015.05.044>.
- Bergin, I.L., Witzmann, F.A., 2013. Nanoparticle toxicity by the gastrointestinal route: evidence and knowledge gaps. *Int. J. Biomed. Nanosci. Nanotechnol.* 3, 1–2. <http://dx.doi.org/10.1504/IJBNN.2013.054515>.
- Bravo-Alvarez, H., Torres-Jardón, R., 2002. Air pollution levels and trends in the México City metropolitan area." Chapter 6. In: urban air pollution and forest: resources at risk in the Mexico City air. In: Basin, M.F., de Bauer, L., Hernández, T. (Eds.), *Ecological Studies* 156. Springer-Verlag, New York, pp. 121–159.
- Brickman, A.M., Gúzman, V.A., Gonzalez-Castellon, M., Razlighi, Q., Gu, Y., Nar-khede, A., Janicki, S., Ichise, M., Stern, Y., Manly, J.J., Schupf, N., Marshall, R.S., 2015. Cerebral autoregulation, beta-amyloid and white matter hyperintensities are interrelated. *Neurosci. Lett.* 592, 54–58.
- Cabezas, R., Avila, M., González, J., El-Bachá, R.S., Báez, E., García-Segura, L.M., Coronel, J.C., Capani, F., Cardona-Gómez, G.P., Barreto, G.E., 2014. Astrocytic modulation of blood brain barrier: perspectives on Parkinson's disease. *Front. Cell Neurosci.* 8, 211. <http://dx.doi.org/10.3389/fncel.2014.0021>.
- Calderón-Garcidueñas, L., Franco-Lira, M., Mora-Tiscareño, A., Medina-Cortina, H., Torres-Jardón, R., Kavanaugh, M., 2013a. Early Alzheimer's and Parkinson's disease pathology in urban children: friend versus foe responses – it is time to face the evidence. *Biomed. Res. Int.* 2013, 161687. <http://dx.doi.org/10.1155/2013/161687>.
- Calderón-Garcidueñas, L., Franco-Lira, M., Rodríguez-Díaz, J., Chao, C., Tompson, C., Mukherjee, P.S., 2015a. Perry CSF biomarkers: low amyloid- $\beta$  1-42 and BDNF and high IFN  $\gamma$  differentiate children exposed to Mexico City high air pollution v controls. *Alzheimer's Disease Uncertainties. J. Alzheimer Dis.* 5, 2. <http://dx.doi.org/10.4172/2161-0460.1000189>.
- Calderón-Garcidueñas, L., Mora-Tiscareño, A., Franco-Lira, M., Zhu, H., Lu, Z., Solorio, E., Torres-Jardón, R., D'Angiulli, A., 2015b. Decreases in Short Term Memory, IQ, and Altered Brain Metabolic Ratios in Urban Apolipoprotein  $\epsilon$ 4 Children Exposed to Air Pollution. *J. Alzheimers Dis.* 45, 757–770.
- Calderón-Garcidueñas, L., Valencia-Salazar, G., Rodríguez-Alcaraz, A., Gambling, T. M., García, R., Osnaya, N., Villarreal-Calderón, A., Devlin, R.B., Carson, J.L., 2001. Ultrastructural nasal pathology in children chronically and sequentially exposed to air pollutants. *Am. J. Respir. Cell. Mol. Biol.* 24, 132–138.
- Calderón-Garcidueñas, L., Mora-Tiscareño, A., Melo-Sánchez, G., Rodríguez-Díaz, J., Torres-Jardón, R., Styner, M., Mukherjee, P.S., Lin, W., Jewells, V., 2015c. A critical proton MR spectroscopy marker of Alzheimer early neurodegenerative change: low hippocampal NAA/Cr ratio impacts APOE 4 Mexico City children and their parents. *J. Alzheimers Dis.* 48, 1065–1075.
- Calderón-Garcidueñas, L., Azzarelli, B., Acuña, H., García, R., Gambling, T.M., Monroy, S., Tizapantzi, M.R., Carson, J.L., Villarreal-Calderon, A., Rewcastle, B., 2002. Air pollution and brain damage. *Toxicol. Pathol.* 30, 373–389.
- Calderón-Garcidueñas, L., Vojdani, A., Blaurock-Busch, E., Busch, I., Friedle, A., Franco-Lira, M., Sarathi-Mukherjee, P., Park, S.B., Torres-Jardón, R., D'Angiulli, A., 2015d. Air pollution and children: neural and tight junction antibodies and combustion metals, the role of barrier breakdown and brain immunity in neurodegeneration. *J. Alzheimers Dis.* 43, 1039–1058.
- Calderón-Garcidueñas, L., Villarreal-Calderon, R., Valencia-Salazar, G., Henríquez-Roldán, C., Gutiérrez-Castrellón, P., Torres-Jardón, R., Osnaya-Brizuela, N., Romero, L., Torres-Jardón, R., Solt, A., Reed, W., 2008c. Systemic inflammation, endothelial dysfunction, and activation in clinically healthy children exposed to air pollutants. *Inhal. Toxicol.* 20, 499–506.
- Calderón-Garcidueñas, L., Mora-Tiscareño, A., Styner, M., Gómez-Garza, G., Zhu, H., Torres-Jardón, R., Carlos, E., Solorio-López, E., Medina-Cortina, H., Kavanaugh, M., D'Angiulli, A., 2012b. White matter hyperintensities, systemic inflammation, brain growth, and cognitive functions in children exposed to air pollution. *J. Alzheimers Dis.* 31, 183–191.
- Calderón-Garcidueñas, L., Serrano-Sierra, A., Torres-Jardón, R., Zhu, H., Yuan, Y., Smith, D., Delgado-Chávez, R., Cross, J.V., Medina-Cortina, H., Kavanaugh, M., Guilarte, T.R., 2013b. The impact of environmental metals in young urbanites' brains. *Exp. Toxicol. Pathol.* 65, 503–511.
- Calderón-Garcidueñas, L., Franco-Lira, M., D'Angiulli, A., Rodríguez-Díaz, J., Blaurock-Busch, E., Busch, I., Chao, C.K., Thompson, C., Mukherjee, P.S., Torres-Jardón, R., Perry, G., 2015e. Mexico City normal weight children exposed to high concentrations of ambient PM<sub>2.5</sub> show high blood leptin and endothelin-1, vitamin D deficiency, and food reward hormone dysregulation versus low pollution controls. Relevance for obesity and Alzheimer disease. *Environ. Res.* 140, 579–592.
- Calderón-Garcidueñas, L., Maronpot, R.R., Torres-Jardón, R., Henríquez-Roldán, C., Schoonhoven, R., Acuña-Ayala, H., Villarreal-Calderon, A., Nakamura, J., Fernando, R., Reed, W., Azzarelli, B., Sweniger, J.A., 2003. DNA damage in nasal and brain tissues of canines exposed to air pollutants is associated with evidence of chronic brain inflammation and neurodegeneration. *Toxicol. Pathol.* 31, 524–538.
- Calderón-Garcidueñas, L., Macías-Parra, M., Hoffmann, H.J., Valencia-Salazar, G., Henríquez-Roldán, C., Monte, O.C., Barragán-Mejía, G., Villarreal-Calderon, R., Romero, L., Granada-Macías, M., Torres-Jardón, R., Medina-Cortina, H., Maronpot, R.R., 2009b. Immunotoxicity and Environment: Immunodysregulation and Systemic Inflammation in Children. *Toxicol. Pathol.* 37, 161–169.
- Calderón-Garcidueñas, L., D'Angiulli, A., Kulesza, R.J., Torres-Jardón, R., Osnaya, N., Romero, L., Keefe, S., Herritt, L., Brooks, D.M., Avila-Ramirez, J., Delgado-Chávez, R., Medina-Cortina, H., González-González, L.O., 2011a. Air pollution is associated with brainstem auditory nuclei pathology and delayed brainstem auditory evoked potentials. *Int. J. Dev. Neurosci.* 29, 365–375.
- Calderón-Garcidueñas, L., Engle, R., Mora-Tiscareño, A., Styner, M., Gómez-Garza, G., Zhu, H., Jewells, V., Torres-Jardón, R., Romero, L., Monroy-Acosta, M.E., Bryant, C., González-González, L.O., Medina-Cortina, H., D'Angiulli, A., 2011b. Exposure to severe urban pollution influences cognitive outcomes, brain volume and systemic inflammation in clinically healthy children. *Brain Cogn.* 77, 345–355.
- Calderón-Garcidueñas, L., Solt, A.C., Henríquez-Roldán, C., Torres-Jardón, R., Nuse, B., Herritt, L., Villarreal-Calderón, R., Osnaya, N., Stone, I., García, R., Brooks, D. M., González-Maciél, A., Reynoso-Robles, R., Delgado-Chávez, R., Reed, W., 2008a. Long-term air pollution exposure is associated with neuroinflammation, an altered innate immune response, disruption of the blood-brain-barrier, ultrafine particulate deposition, and accumulation of amyloid beta-42 and alpha-synuclein in children and young adults. *Toxicol. Pathol.* 36, 289–310.
- Calderón-Garcidueñas, L., Kavanaugh, M., Block, M.L., D'Angiulli, A., Delgado-Chávez, R., Torres-Jardón, R., González-Maciél, A., Reynoso-Robles, R., Osnaya, N., Villarreal-Calderon, R., Guo, R., Hua, Z., Zhu, H., Perry, G., Neuroinflammation, Diaz P., 2012a. hyperphosphorylated tau, diffuse amyloid plaques and down-regulation of the cellular prion protein in air pollution exposed children and adults. *J. Alzheimer Dis.* 28, 93–107.
- Calderón-Garcidueñas, L., Vincent, R., Mora-Tiscareño, A., Franco-Lira, M., Henríquez-Roldán, C., Barragán-Mejía, G., Garrido-García, L., Camacho-Reyes, L., Valencia-Salazar, G., Paredes, R., Romero, L., Osnaya, H., Villarreal-Calderón, R., Torres-Jardón, R., Hazucha, M.J., Reed, W., 2007. Elevated plasma endothelin-1 and pulmonary arterial pressure in children exposed to air pollution. *Environ. Health Perspect.* 115, 1248–1253.
- Calderón-Garcidueñas, L., Franco-Lira, M., Henríquez-Roldán, C., González-Maciél, A., Reynoso-Robles, R., Villarreal-Calderon, R., Herritt, L., Brooks, D., Keefe, S., Palacios-Moreno, J., Villarreal-Calderon, R., Torres-Jardón, R., Medina-Cortina, H., Delgado-Chávez, R., Aiello-Mora, M., Maronpot, R.R., Doty, R.L., 2010. Urban air pollution: influences on olfactory function and pathology in exposed children and young adults. *Exp. Toxicol. Pathol.* 62, 91–102.
- Calderón-Garcidueñas, L., Mora-Tiscareño, A., Gómez-Garza, G., Carrasco-Portugal, M., del, C., Pérez-Guillé, B., Flores-Murrieta, F.J., Pérez-Guillé, G., Juarez-Olguin, H., Monroy, M.E., Monroy, S., González-Maciél, A., Reynoso-Robles, R., Villarreal-Calderón, R., Patel, S.A., Kumarathanan, P., Vincent, R., Henríquez-Roldán, C., Torres-Jardón, R., Maronpot, R., 2009a. Effects of a cyclooxygenase 2 preferential inhibitor in young healthy dogs exposed to air pollution: a pilot study. *Toxicol. Pathol.* 37, 644–660.
- Calderón-Garcidueñas, L., Mora-Tiscareño, A., Ontiveros, E., Gómez-Garza, G., Barragán-Mejía, G., Broadway, J., Chapman, S., Valencia-Salazar, G., Jewells, V., G., Maronpot, R.R., Henríquez-Roldán, C., Pérez-Guillé, B., Torres-Jardón, R., Herritt,

- L., Brooks, D., Osnaya-Brizuela, N., Monroy, M.E., González-Maciel, A., Reynoso-Robles, R., Villarreal-Calderon, R., Solt, A.C., Engle, R.W., 2008b. Air pollution, cognitive deficits and brain abnormalities: A pilot study with children and dogs. *Brain Cogn.* 68, 117–127.
- Carson, J.L., Brighton, L.E., Collier, A.M., Bromberg, P.A., 2013. Correlative ultrastructural investigations of airway epithelium following experimental exposure to defined air pollutants and lifestyle exposure to tobacco smoke. *Inhal. Toxicol.* 25, 134–140.
- Castejón, O.J., 2012. Ultrastructural pathology of endothelial tight junctions in human brain edema. *Folia Neuropathol.* 50, 118–129.
- Castejón, O.J., 2014. Ultrastructural alterations of human cortical capillary basement membrane in human brain oedema. *Folia Neuropathol.* 52, 10–21.
- Diamond, B., Honig, G., Mader, S., Brimberg, L., Volpe, B.T., 2014. Brain-reactive antibodies and disease. *Ann. Rev. Immunol.* 31, 345–385.
- Edwards, J.D., Ramírez, J., Black, S.E., 2015. Unraveling the potential co-contributors of cerebral small vessel vasculopathy to the pathogenesis of Alzheimer's dementia. *Alzheimers Res. Ther.* 7 (1), 49. <http://dx.doi.org/10.1186/s13195-015-0133-2>.
- Eisenhut, M., 2014. Vasospasm in cerebral inflammation. *Int. J. Inflamm.* 2014, 509707. <http://dx.doi.org/10.1155/2014/509707>.
- Elahy, M., Jackaman, C., Mamo, J.C., Lam, V., Dhaliwal, S.S., Giles, C., Nelson, D., Takechi, R., 2015. Blood-brain-barrier dysfunction developed during normal aging is associated with inflammation and loss of tight junctions but not with leukocyte recruitment. *Immun. Ageing* 12, 2. <http://dx.doi.org/10.1186/s12979-015-0029-9>.
- Finch, J., Conklin, D.J., 2015. Air pollution-induced vascular dysfunction: potential role of endothelin-1 (ET-1) system. *Cardiovasc. Toxicol.*
- Fischer, F.U., Wolf, D., Scheurich, A., Fellgiebel, A., 2015. Alzheimer's disease neuroimaging initiative. Altered whole-brain white matter networks in preclinical Alzheimer's disease. *Neuroimage Clin.* 8, 660–666. <http://dx.doi.org/10.1016/j.nicl.2015.06.007>.
- Garwood, C.J., Simpson, J.E., Al Mashhadi, S., Axe, C., Wilson, S., Heath, P.R., Shaw, P., Matthews, F.E., Brayne, C., Ince, P.G., Wharton, S.B., 2014. MRC cognitive function and ageing Study. DNA damage response and senescence in endothelial cells of human cerebral cortex and relation to Alzheimer's neuropathology progression: a population-based study in the Medical Research Council Cognitive Function and Ageing Study (MRC-CFAS) cohort. *Neuropathol. Appl. Neurobiol.* 40, 802–814.
- Gehr, P., Clift, M.J., Brandenberger, C., Lehmann, A., Herzog, F., Rothen-Rutishauser, B., 2011. Endocytosis of environmental and engineered micro- and nanosized particles. *Comp. Physiol.* 1, 1159–1174.
- Grade, M., Hernández-Tamames, J.A., Pizzini, F.B., Achten, E., Golay, X., Smits, M., 2015. A neuroradiologist's guide to arterial spin labeling MRI in clinical practice. *Neuroradiology* 57, 1181–1202, Sep 9.
- Haight, T.J., Bryan, R.N., Erus, G., Davatzikos, C., Jacobs, D.R., D'Esposito, M., Lewis, C.E., Launer, L.J., 2015. Vascular risk factors, cerebrovascular reactivity and the default-mode brain network. *Neuroimage* 115, 7–16. <http://dx.doi.org/10.1016/j.neuroimage.2015.04.039>.
- Harkema, J.R., Carey, S.A., Wagner, J.G., 2006. The nose revisited: a brief review of the comparative structure, function, and toxicologic pathology of the nasal epithelium. *Toxicol. Pathol.* 34, 252–269.
- Haseloff, R.F., Dithmer, S., Winkler, L., Wolburg, H., Blasig, I.E., 2015. Transmembrane proteins of the tight junctions at the blood-brain barrier: structural and functional aspects. *Semin. Cell Dev. Biol.* 38, 16–25. <http://dx.doi.org/10.1016/j.semcdb.2014.11.004>.
- Hawkes, C.A., Gentleman, S.M., Nicoll, J.A., Carare, R.O., 2015. Prenatal high-fat diet alters the cerebrovasculature and clearance of  $\beta$ -amyloid in adult offspring. *J. Pathol.* 235, 619–631.
- Hawkes, C.A., Jayakody, N., Johnston, D.A., Bechmann, I., Carare, R.O., 2014. Failure of perivascular drainage of  $\beta$ -amyloid in cerebral amyloid angiopathy. *Brain Pathol.* 24, 396–403.
- Hung, V.K., Yeung, P.K., Lai, A.K., Ho, M.C., Chan, K.C., Wu, E.X., Chung, S.S., Cheung, C.W., Chung, S.K., 2015. Selective astrocytic endothelin-1 overexpression contributes to dementia associated with ischemic stroke by exaggerating astrocyte-derived amyloid secretion. *J. Cereb. Blood Flow. Metab.* 35, 1687–1696. <http://dx.doi.org/10.1038/jcbfm.2015.109>.
- Iadecola, C., 2004. Neurovascular regulation in the normal brain and in Alzheimer's disease. *Nat. Rev. Neurosci.* 5, 347–360.
- Jung, C.R., Lin, Y.T., Hwang, B.F., 2015. Ozone particulate matter and newly diagnosed Alzheimer's disease: a population-based cohort study in Taiwan. *J. Alzheimers Dis.* 44, 573–584.
- Kamat, P.K., Swarnkar, S., Rai, S., Kumar, V., Tyagi, N., 2014. Astrocyte mediated MMP-9 activation in the synapse dysfunction: an implication in Alzheimer disease. *Ther. Targets Neurol. Dis.* 1 (1), e243, pii.
- Kaplan, G.G., Hubbard, J., Korzenik, J., Sands, B.E., Panaccione, R., Ghosh, S., Wheeler, A.J., Villeneuve, P.J., 2010. The inflammatory bowel diseases and ambient air pollution: a novel association. *Am. J. Gastroenterol.* 105, 2412–2419.
- Karmakar, A., Zhang, Q., Zhang, Y., 2014. Neurotoxicity of nanoscale materials. *J. Food Drug Anal.* 22, 147–160.
- Karthikeyan, S., Thomson, E.M., Kumarathasan, P., Guénette, J., Rosenblatt, D., Chan, T., Rideout, G., Vincent, R., 2013. Nitrogen dioxide and ultrafine particles dominate the biological effects of inhaled diesel exhaust treated by a catalyzed diesel particulate filter. *Toxicol. Sci.* 135, 437–450.
- Keaney, J., Campbell, M., 2015. The dynamic blood-brain-barrier. *FEBS J.* 282, 4067–4079. <http://dx.doi.org/10.1111/febs.13412>, Aug 16.
- Lacoste, B., Gu, C., 2015. Control of cerebrovascular patterning by neural activity during postnatal development. *Mech. Dev.* 138, 43–49. <http://dx.doi.org/10.1016/j.mod.2015.06.003>.
- Lai, A.Y., Dorr, A., Thomason, L.A., Koletar, M.M., Sled, J.G., Stefanovic, B., McLaurin, J., 2015. Venular degeneration leads to vascular dysfunction in a transgenic model of Alzheimer's disease. *Brain* 138 (Pt 4), 1046–1058. <http://dx.doi.org/10.1093/brain/awv023>.
- Laman, J.D., Weller, R.O., 2013. Drainage of cells and soluble antigen from the CNS to regional lymph nodes. *J. Neuroimmune Pharmacol.* 8, 840–856.
- Lecrux, C., Hamel, E., 2011. The neurovascular unit in brain function and disease. *Acta Physiol.* 203, 47–59.
- Lei, Q., Li, S., Zheng, R., Xu, K., Li, S., 2015. Endothelin-1 expression and alterations of cerebral microcirculation after experimental subarachnoid hemorrhage. *Neuroradiology* 57, 63–70.
- Levin, E.C., Acharya, N.K., Han, M., Zavareh, S.B., Sedeyn, J.C., Venkataraman, V., Nagele, R.G., 2010. Brain-reactive autoantibodies are nearly ubiquitous in human sera and may be linked to pathology in the context of blood-brain barrier breakdown. *Brain Res.* 1345, 221–232.
- Li, C.H., Jhan, C., Cheng, Y.W., Tsai, C.H., Liu, C.W., Lee, C.C., Chen, R.M., Shyu, M.K., Kang, J.J., 2015. Gold nanoparticles increase endothelial paracellular permeability by altering components of endothelial tight junctions, and increase blood-brain barrier permeability in mice. *Toxicol. Sci.* 148, 192–203, Aug 13. pii: kfv176.
- Licht, T., Keshet, E., 2015. The vascular niche in adult neurogenesis. *Mech. Dev.* 138, 56–62. <http://dx.doi.org/10.1016/j.mod.2015.06.001>, pii: S0925-4773(15)00045-3.
- Ljubimova, J.Y., Kleinman, M.T., Karabalin, N.M., Inoue, S., Konda, B., Gangalum, P., Markman, J.L., Ljubimov, A.V., Black, K.L., 2013. Gene expression changes in rat brain after short and long exposures to particulate matter in Los Angeles basin air. Comparison with human brain tumors. *Exp. Toxicol. Pathol.* 65, 1063–1071.
- Lo, E.H., Rosenberg, G.A., 2009. The neurovascular unit in health and disease. *Stroke* 40, S2–S3.
- Lourenco, C.F., Ledo, A., Dias, C., Barbosa, R.M., Laranjinha, J., 2015. Neurovascular and neurometabolic derailment in aging and Alzheimer's disease. *Front. Aging Neurosci.* 7, 103. <http://dx.doi.org/10.3389/fnagi.2015.00103>.
- Love, S., Miners, J.S., 2015. White matter hypoperfusion and damage in dementia: post-mortem assessment. *Brain Pathol.* 25, 99–107.
- Mugica, V., Torres, M., Salinas, E., Gutiérrez, M., García, R., 2010. Polycyclic aromatic hydrocarbons in the urban atmosphere of Mexico City. In: Villanyi, Vanda (Ed.), *Air Pollution*. InTech, Croatia, pp. 73–97. <http://www.intechopen.com/books/air-pollution/polycyclic-aromatic-hydrocarbons-in-the-urban-atmosphere-of-mexico-city>.
- Maillard, P., Fletcher, E., Harvey, D., Carmichael, O., Reed, B., Mungas, D., DeCarli, C., 2011. White matter hyperintensity penumbra. *Stroke* 42, 1917–1922.
- Mathiisen, T.M., Lehre, K.P., Danbolt, N.C., Ottersen, O.P., 2010. The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia* 58, 1094–1103. <http://dx.doi.org/10.1002/glia.20990>.
- Meng, N., Han, L., Pan, X., Su, L., Jiang, Z., Lin, Z., Zhao, J., Zhang, S., Zhang, Y., Zhao, B., Miao, J., 2015. Nano-Mg(OH)<sub>2</sub>-induced proliferation inhibition and dysfunction of human umbilical vein vascular endothelial cells through caveolin-1-mediated endocytosis. *Cell Biol. Toxicol.* 31, 15–27.
- Molina, L.T., Madronich, S.J., Gaffney, J.S., Apel, E., de Foy, B., Fast, J., Ferrare, R., Herndon, S., Jimenez, J.L., Lamb, B., Osornio-Vargas, A.R., Russell, P., Schauer, J.J., Stevens, P.S., Volkamer, R., Zavala, M., 2010. An overview of the MILAGRO 2006 Campaign: Mexico City emissions and their transport and transformation. *Atmos. Chem. Phys.* 10, 8697–8760.
- Morris, A.W., Carare, R.O., Schreiber, S., Hawkes, C.A., 2014. The cerebrovascular basement membrane: role in the clearance of  $\beta$ -amyloid and cerebral amyloid angiopathy. *Front. Aging Neurosci.* 19, 251. <http://dx.doi.org/10.3389/fnagi.2014.00251>.
- Muir, A.R., Peter, A., 1962. Quintuple-layered membrane junctions at terminal bars between endothelial cells. *J. Cell Biol.* 12, 443–448.
- Murray, K.N., Girard, S., Holmes, W.M., Parkes, L.M., Williams, S.R., Parry-Jones, A.R., Allan, S.M., 2014. Systemic inflammation impairs tissue reperfusion through endothelin-dependent mechanisms in cerebral ischemia. *Stroke* 45, 3412–3419.
- Nguemni, C., Gomez-Smith, M., Jeffers, M.S., Schuch, C.P., Corbett, D., 2015. Time course of neuronal death following endothelin-1 induced focal ischemia in rats. *J. Neurosci. Methods* 242, 72–76.
- Nir, T.M., Jahanshad, N., Villalon-Reina, J.E., Toga, A.W., Jack, C.R., Weiner, M.W., Thompson, P.M., 2013. Alzheimer's Disease Neuroimaging Initiative (ADNI). Effectiveness of regional DTI measures in distinguishing Alzheimer's disease, MCI and normal aging. *Neuroimage Clin.* 180–195.
- Prins, N.D., Scheltens, P., 2015. White matter hyperintensities, cognitive impairment and dementia: an update. *Nat. Rev. Neurol.* 11, 157–165.
- Promjunyakul, N., Lahna, D., Kaye, J.A., Dodge, H.H., Erten-Lyons, D., Rooney, W.D., Silbert, L.C., 2015. Characterizing the white matter hyperintensity penumbra with cerebral blood flow measures. *Neuroimage Clin.* 8, 224–229.
- Qosa, H., Miller, D.S., Pasinelli, P., Trotti, D., 2015. Regulation of ABC efflux transporters at blood-brain barrier in health and neurological disorders. *Brain Res.* 1628, 298–316, Jul 15. pii: S0006-8993(15)00530-2.
- Querol, X., Pey, J., Minguillón, M.C., Pérez, N., Alastuey, A., Viana, M., Moreno, T., Bernabé, R.M., Blanco, S., Cárdenas, B., Vega, E., Sosa, G., Escalona, S., Ruiz, H., Artífano, B.P.M., 2008. Speciation and sources in Mexico during the MILAGRO-2006 Campaign. *Atmos. Chem. Phys.* 8, 111–128.
- Retama, A., Baumgardner, D., Raga, G.B., McMeeking, G.R., Walker, J.W., 2015. Seasonal and diurnal trends in black carbon properties and co-pollutants in

- Mexico City. *Atmos. Chem. Phys.* 15, 9693–9709.
- Ritchie, S.J., Bastin, M.E., Tucker-Drob, E.M., Maniega, S.M., Engelhardt, L.E., Cox, S.R., Royle, N.A., Gow, A.J., Corley, J., Pattie, A., Taylor, A.M., Valdés Hernández, M., del, C., Starr, J.M., Wardlaw, J.M., Deary, I.J., 2015. Coupled changes in brain white matter microstructure and fluid intelligence in later life. *J. Neurosci.* 35, 8672–8682.
- Seinfeld, J.H., Pandis, S.N., 1998. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 1st edition. J. Wiley, New York.
- Sharma, A., Muresanu, D.F., Patnaik, R., Sharma, H.S., 2013. Size and age-dependent neurotoxicity of engineered metal nanoparticles in rats. *Mol. Neurobiol.* 48, 386–396.
- Simard, M., Arcuino, G., Takano, T., Liu, Q.S., Nedergaard, M., 2003. Signaling at the gliovascular surface. *J. Neurosci.* 23, 9254–9262.
- Spigoni, V., Cito, M., Alinovi, R., Pinelli, S., Passeri, G., Zavaroni, I., Goldoni, M., Campanini, M., Aliatis, I., Mutti, A., Bonadonna, R.C., Dei Cas, A., 2015. Effects of TiO<sub>2</sub> and CO<sub>2</sub>O<sub>4</sub> nanoparticles on circulating angiogenic cells. *PLoS One* 10, e0119310. <http://dx.doi.org/10.1371/journal.pone.0119310>.
- Sweeney, M.D., Sagare, A.P., Zlokovic, B.V., 2015. Cerebrospinal fluid biomarkers of neurovascular dysfunction in mild dementia and Alzheimer's disease. *J. Cereb. Blood Flow Metab.* 35, 1055–1068.
- Thompson, E.A., Sayers, B.C., Glista-Baker, E.E., Shipkowski, K.A., Taylor, A.J., Bonner, J.C., 2014. Innate immune responses to nanoparticle exposure in the lung. *J. Environ. Immunol. Toxicol.* 1, 150–156.
- Thomson, E.M., Kumarathasan, P., Goegan, P., Aubin, R.A., Vincent, R., 2005. Differential regulation of the lung endothelin system by urban particulate matter and ozone. *Toxicol. Sci.* 88, 103–113.
- Tietz, S., Engelhardt, B., 2015. Brain barriers: Crosstalk between complex tight junctions and adherens junctions. *J. Cell Biol.* 209, 493–506. <http://dx.doi.org/10.1083/jcb.201412147>.
- Ucciferri, N., Collnot, E.M., Gaiser, B.K., Tirella, A., Stone, V., Domenici, C., Lehr, C.M., Ahluwalia, A., 2014. In vitro toxicological screening of nanoparticles on primary human endothelial cells and the role of flow in modulating cell response. *Nanotoxicology* 8, 697–708.
- Van Miert, E., Dumont, X., Bernard, A., 2005. CC16 as a marker of lung epithelial hyperpermeability in an acute model of rats exposed to mainstream cigarette smoke. *Toxicol. Lett.* 159, 115–123.
- Vega, E., Eidels, S., Ruiz, H., López-Veneroni, D., Sosa, G., González, E., Gasca, J., Mora, V., Reyes, E., Sánchez-Reyna, G., Villaseñor, R., Chow, J.C., Watson, J.G., Edgerton, S.A., 2010. Particulate air pollution in Mexico City: A detailed view. *Aerosol Air Qual. Res.* 10, 193–211.
- Villarreal-Calderón, A., Acuña, H., Villarreal-Calderón, J., Garduño, M., Henríquez-Roldán, C.F., Calderón-Garcidueñas, L., Valencia-Salazar, G., 2002. Assessment of physical education time and after-school outdoor time in elementary and middle school students in south Mexico City: the dilemma between physical fitness and the adverse health effects of outdoor pollutant exposure. *Arch. Environ. Health* 57, 450–460.
- Wang, Y., Jin, S., Sonobe, Y., Cheng, Y., Horiuchi, H., Parajuli, B., Kawanokuchi, J., Mizuno, T., Takeuchi, H., Suzumura, A., 2014. Interleukin-1 $\beta$  induces blood-brain-barrier disruption by downregulating Sonic hedgehog in astrocytes. *PLoS One* 9, e110024. <http://dx.doi.org/10.1371/journal.pone.0110024>.
- Wang, Z., Tirupathi, C., Minshall, R.D., Malik, A.B., 2009. Size and dynamics of caveolae studied using nanoparticles in living endothelial cells. *ACS Nano* 3, 4110–4116.
- Wardlaw, J.M., Smith, E.E., Biessels, G.H., Cordonnier, C., Fazekas, F., Frayne, R., Lindley, R.I., O'Brien, J.T., Barkhof, F., Benavente, O.R., Black, S.E., Brayne, C., Breteler, M., Chabriat, H., DeCarli, C., de Leeuw, F.E., Doubal, F., Duering, M., Fox, N.C., Greenberg, S., Hachinski, V., Kilimann, I., Mok, V., van Oostenbrugge, R., Pantoni, L., Speck, O., Stephan, B.C.M., Teipel, S., Viswanathan, A., Werring, D., Chen, C., Smith, C., van Buchem, M., Norrving, B., Gorelick, P.B., Dichgans, M., 2013. Standards for reporting vascular changes on nEuroimaging (STRIVE v1). Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *Lancet Neurol.* 12, 822–838.
- Weinl, C., Castaneda-Vega, S., Riehle, H., Stritt, C., Calaminus, C., Wolburg, H., Muel, S., Breithaupt, A., Gruber, A.D., Wasyluk, B., Olson, E.N., Adams, R.H., Pichler, B.J., Nordheim, A., 2015. Endothelial depletion of murine SRF/MRTF provokes intracerebral hemorrhagic stroke. *Proc. Natl. Acad. Sci. USA* 112, 9914–9919.
- Weller, R.O., Boche, D., Nicoll, J.A., 2009. Microvasculature changes and cerebral amyloid angiopathy in Alzheimer's disease and their potential impact on therapy. *Acta Neuropathol.* 118, 87–102.
- Weller, R.O., Hawkes, C.A., Kalaria, R.N., Werring, D.J., Carare, R.O., 2015. White matter changes in dementia: role of impaired drainage of interstitial fluid. *Brain Pathol.* 25, 63–78.
- Winkler, E.A., Sagare, A.P., Zlokovic, B.V., 2014. The pericyte: a forgotten cell type with important implications for Alzheimer's disease? *Brain Pathol.* 24, 371–386. <http://dx.doi.org/10.1111/bpa.12152>.
- Wyss-Coray, T., Lin, C., Sanan, D.A., Mucke, L., Masliah, E., 2000. Chronic overproduction of Transforming growth factor  $\beta$ -1 by astrocytes promotes Alzheimer's disease-like microvascular degeneration in transgenic mice. *Am. J. Pathol.* 156, 139–150.
- Yu, S.H., Tang, D.W., Hsieh, H.Y., Wu, W.S., Lin, B.X., Chuang, E.Y., Sung, H.W., Mi, F.L., 2013. Nanoparticle-induced tight-junction opening for the transport of an anti-angiogenic sulfated polysaccharide across Caco-2 cell monolayers. *Acta Biomater.* 9, 7449–7459.
- Zlokovic, B.V., 2008. The blood-brain-barrier in health and chronic neurodegenerative disorders. *Neuron* 57, 178–201.